

08-14-00 Seq. A

08/11/00 09/637302 08/11/00

jc901 U.S. PTO

TRANSMITTAL OF UTILITY PATENT APPLICATION FOR FILING

Certification under 37 CFR 1.10 (if applicable)

EL154221135US

August 11, 2000

"Express Mail" mailing label number

Date of Deposit

I hereby certify that this Transmittal letter, enclosed application, and any other documents referred to as enclosed herein, are being deposited in an envelope with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.

Cedric Rodgers

(Typed or printed name of person mailing application)

Cedric Rodgers

(Signature of person mailing application)

ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
Washington, D.C. 20231

Attorney Docket No. TSRI 710.2

Customer Number 002387

Sir:

Transmitted herewith for filing is the utility patent application of inventor(s): John Hood, Brian Eliceiri and David A. Cheresch.

and entitled: **METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS
USING TYROSINE KINASE RAF AND RAS**

1. Type Of Application

This application is:

- ☐ an original (nonprovisional) application.
☐ a divisional of prior application Serial No. _____.
☐ a continuation of prior application Serial No. _____.
☐ a continuation-in-part of prior application Serial No. _____.
☒ an application claiming priority of U.S. Provisional Application Nos. 60/148,924 and 60/215,951

☐ The entire disclosure of the prior application is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

☐ The prior application is assigned of record to: _____

☐ Additional prior application information: Examiner _____ Group _____

2. Enclosed Application Elements are:

- ☒ A duplicate copy of this transmittal letter,
☒ specification (including claims and abstract) containing 60 pages,
☒ drawings: ☐ 1 copy of _____ sheet(s) of formal drawings, OR
☒ 1 copy of Twenty (20) sheet(s) of informal drawings. (Figs. 1-20B)
☐ an executed declaration or oath for the utility patent application including a power of attorney, OR
☒ an unexecuted declaration or oath for the utility patent application including a power of attorney, OR
☐ a copy of an executed declaration or oath including power of attorney from a priority application,
☐ statement deleting inventor(s) named in the priority application
☐ Microfiche Computer Program
☒ nucleotide and/or amino acid sequence
a. ☒ computer readable copy
b. ☒ paper copy
c. ☒ statement verifying above copies

TRANSMITTAL OF UTILITY PATENT APPLICATION FOR FILING
Page 2

3. Enclosed Accompanying Application Parts are:

- ☐ One (1) Verified Statement(s) relating to small entity status.
- ☐ Copy of verified statement filed in prior application; status still proper and desired.
- ☐ Preliminary Amendment
- ☐ Claim cancellations are indicated in Preliminary Amendment
- ☒ one itemized, stamped, and self-addressed postcard for the PTO Mail Room date stamp.
- ☐ English translation document
- ☐ Information Disclosure Statement including Form PTO-1449 and copies of the citations therein.

4. Filing Fees (as calculated below)

(Col. 1)		(Col. 2)		
For:	Number Filed	Number Extra	Rate	Fee
Basic Fee				\$ 690
Total Claims	67 — 20	47	x \$ 18 =	\$ 846
Independent Claims	13 - 3	10	x \$ 78 =	\$ 780
Multiple Dependent Claim Presented (if applicable)			+ \$260 =	\$ N/A
Subtotal				\$ 2,316
Reduction by 50% for filing by small entity				\$ -0-
* If the difference in Col. 1 is less than zero, enter "0" in Col. 2.				
TOTAL				\$ 2,316

- ☐ Please charge my Deposit Account No. 15-0508 in the amount of \$_____.
- ☒ A check in the amount of \$2,316.00 to cover the filing fee is enclosed.
- ☒ The Commissioner is authorized to charge payment of the following amounts associated with this communication or credit any overpayment to Deposit Account No. 15-0508:
- ☒ Additional filing fees under 37 CFR 1.16 or deficiencies in remittances therefor.
- ☒ Additional processing fees under 37 CFR 1.17 or deficiencies in remittances therefor.
- ☒ **ONLY if applicant has partially paid** the patent issue fee under 37 C.F.R. §1.18, then the **deficiency** shall be charged to Deposit Account No. 15-0508, and the Commissioner is authorized to so charge the Deposit Account.
- ☒ The Commissioner is hereby generally authorized under 37 CFR 1.136(a)(3) to treat any future reply in this or any related application filed pursuant to 37 CFR 1.53 requiring an extension of time as incorporating a request therefor, and the Commissioner is hereby specifically authorized to charge Deposit Account No. 15-0508 for any fee that may be due in connection with such a request for an extension of time.

Date: August 11, 2000

Attorney's Signature



Mark Chao (Reg. No. 37,293)

Correspondence Address:

Mark Chao (Reg. No. 37,293)
OLSON & HIERL, LTD.
20 North Wacker Drive, 36th Floor
Chicago, Illinois 60606
Telephone: (312) 580-1180

METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS USING PROTEIN KINASE RAF AND RAS

Cross-reference to Related Applications

This application claims priority to U.S. Provisional Patent Application Serial
5 No. 60/215,951 filed July 5, 2000, and United States Provisional Patent Application
Serial No. 60/148,924, filed August 13, 1999.

Technical Field

The present invention relates generally to the field of medicine, and relates
specifically to methods and compositions for modulating angiogenesis of tissues using
10 the protein kinase Raf or Ras, variants of Raf or Ras, using reagents which modulate
Raf or Ras, and using nucleic acids encoding them.

Background

Angiogenesis is a process of tissue vascularization that involves the growth of
new blood vessels into a tissue, and is also referred to as neo-vascularization. The
15 process is mediated by the infiltration of endothelial cells and smooth muscle cells.
The process is believed to proceed in any one of three ways: the vessels can sprout
from pre-existing vessels, de-novo development of vessels can arise from precursor
cells (vasculogenesis), or existing small vessels can enlarge in diameter. Blood et al.,
Bioch. Biophys. Acta, 1032:89-118 (1990).

20 Angiogenesis is an important process in neonatal growth, but is also important
in wound healing and in the pathogenesis of a large variety of clinical diseases
including tissue inflammation, arthritis, tumor growth, diabetic retinopathy, macular
degeneration by neovascularization of the retina and like conditions. These clinical
manifestations associated with angiogenesis are referred to as angiogenic diseases.
25 Folkman et al., Science, 235:442-447 (1987). Angiogenesis is generally absent in
adult or mature tissues, although it does occur in wound healing and in the corpus
luteum growth cycle. See, for example, Moses et al., Science, 248:1408-1410 (1990).

It has been proposed that inhibition of angiogenesis would be a useful therapy
for restricting tumor growth. Inhibition of angiogenesis has been proposed by (1)

inhibition of release of “angiogenic molecules” such as bFGF (basic fibroblast growth factor), (2) neutralization of angiogenic molecules, such as by use of anti-bFGF antibodies, (3) use of inhibitors of vitronectin receptor $\alpha_v\beta_3$, and (4) inhibition of endothelial cell response to angiogenic stimuli. This latter strategy has received attention, and Folkman et al., Cancer Biology, 3:89-96 (1992), have described several endothelial cell response inhibitors, including collagenase inhibitor, basement membrane turnover inhibitors, angiostatic steroids, fungal-derived angiogenesis inhibitors, platelet factor 4, thrombospondin, arthritis drugs such as D-penicillamine and gold thiomalate, vitamin D₃ analogs, alpha-interferon, and the like that might be used to inhibit angiogenesis. For additional proposed inhibitors of angiogenesis, see Blood et al., Bioch. Biophys. Acta, 1032:89-118 (1990), Moses et al., Science, 248:1408-1410 (1990), Ingber et al., Lab. Invest., 59:44-51 (1988), and United States Patent Nos. 5,092,885, 5,112,946, 5,192,744, 5,202,352, 5,753,230 and 5,766,591. None of the inhibitors of angiogenesis described in the foregoing references involve the Raf proteins, however.

For angiogenesis to occur, endothelial cells must first degrade and cross the blood vessel basement membrane in a manner similar to that used by tumor cells during invasion and metastasis formation.

It has been previously reported that angiogenesis depends on the interaction between vascular integrins and extracellular matrix proteins. Brooks et al., Science, 264:569-571 (1994). Furthermore, it was reported that programmed cell death (apoptosis) of angiogenic vascular cells is initiated by the interaction, which would be inhibited by certain antagonists of the vascular integrin $\alpha_v\beta_3$. Brooks et al., Cell, 79:1157-1164 (1994). More recently, it has been reported that the binding of matrix metalloproteinase-2 (MMP-2) to vitronectin receptor ($\alpha_v\beta_3$) can be inhibited using $\alpha_v\beta_3$ antagonists, and thereby inhibit the enzymatic function of the proteinase. Brooks et al., Cell, 85:683-693 (1996).

Summary of the Invention

The present invention contemplates modulation of angiogenesis in tissues where that angiogenesis depends upon the activity of protein kinase Raf, also referred to generically herein as Raf.

5 Compositions and methods for modulating angiogenesis in a tissue associated with a disease condition are contemplated. A composition comprising an angiogenesis-modulating amount of a Raf protein is administered to tissue to be treated for a disease condition that responds to modulation of angiogenesis. The composition providing the Raf protein can contain purified protein, biologically active
10 protein fragments, recombinantly produced Raf protein or protein fragments or fusion proteins, or gene/nucleic acid expression vectors for expressing a Raf protein.

Where the Raf protein is inactivated or inhibited, the modulation is an inhibition of angiogenesis. Where the Raf protein is active or activated, the modulation is a potentiation of angiogenesis.

15 The tissue to be treated can be any tissue in which modulation of angiogenesis is desirable. For angiogenesis inhibition, it is useful to treat diseased tissue where deleterious neovascularization is occurring. Exemplary tissues include inflamed tissue, solid tumors, metastases, tissues undergoing restenosis, and the like tissues.

For potentiation, it is useful to treat patients with hypoxic tissues such as those
20 following stroke, myocardial infarction or associated with chronic ulcers, tissues in patients with ischemic limbs in which there is abnormal, i.e., poor circulation, due to diabetic or other conditions. Patients with chronic wounds that do not heal, and therefore could benefit from the increase in vascular cell proliferation and neovascularization, can be treated as well.

25 Particularly preferred is the use of Raf protein containing a modified amino acid sequence as described herein. Several particularly useful modified Raf proteins, including Raf fusion proteins such as Raf-caax and nucleic acid constructs which encode for the expression thereof are described herein and are within the purview of the present invention.

The present invention also encompasses a pharmaceutical composition suitable for inhibiting angiogenesis in a target mammalian tissue comprising a viral or non-viral gene transfer vector containing a nucleic acid, the nucleic acid having a nucleic acid segment encoding for a Raf protein, and the Raf protein having any amino acid residue at codon 375 except for lysine, and a pharmaceutically acceptable carrier or excipient. A particularly preferred embodiment utilizes Raf protein designated as Raf K375M and described in the examples below. Another inactive Raf construct is a nucleic acid which encodes for a Raf protein having the carboxy terminal portion deleted. One preferred embodiment utilizes a Raf protein designated Raf 1-305, which is an inactive Raf protein.

Also envisioned is a pharmaceutical composition suitable for stimulating angiogenesis in a target mammalian tissue and comprising a viral or non-viral gene transfer vector containing a nucleic acid having a segment encoding for a Raf protein having kinase activity and a pharmaceutically acceptable carrier or excipient therefor. A preferred nucleic acid encodes for an inhibitory Raf fusion protein that is Raf-caax. Another inhibitory Raf construct contains a nucleic acid encoding for a Raf protein having the amino terminal portion of the protein deleted. One preferred embodiment utilizes a Raf protein designated Raf 306-648, and described in the examples below.

The invention further contemplates modulation of angiogenesis in tissues by small GTPase Ras, also referred to generically herein as Ras, due to its role in signaling Raf, as described herein. Also envisioned is the modulation of angiogenesis in tissues utilizing the combination of Ras and Raf modulation. Such combined modulation can take the form of a single administration of combined formulations of protein, or nucleic acid encoding modulating protein, or the separate administration of individual doses, in an angiogenesis-modulating amount.

Compositions and methods for modulating angiogenesis in a tissue, associated with a disease condition are contemplated, where the modulation is directed to the Raf-mediated angiogenesis pathway via the Ras protein. A composition comprising an angiogenesis-modulating amount of a Ras protein is administered to tissue to be

treated for a disease condition that responds to modulation of angiogenesis. The composition providing the Ras protein can contain purified protein, biologically active Ras protein fragments, recombinantly produced Ras protein or protein fragments or fusion proteins, or gene/nucleic acid expression vectors for expressing a Ras protein.

5 Where the Ras protein is inactivated or inhibited, the modulation is an inhibition of angiogenesis. Where the Ras protein is active or activated, the modulation is a potentiation of angiogenesis. Pharmaceutical compositions and methods of use for dominant negative Ras proteins, such as S17N Ras or V12C40 Ras, are contemplated for use in a manner similar to that for proteins of the Raf family. In
10 a further aspect of this invention, pharmaceutical compositions and methods of use for dominant active Ras proteins, such as G12V Ras or V12S35 Ras, are contemplated for uses comparable to those for the Raf family proteins.

 Further contemplated are methods for modulating angiogenesis in a tissue associated with a disease condition comprising administering an angiogenesis
15 modulating amount of a pharmaceutical composition comprising a Raf protein or a nucleotide sequence capable of expressing Raf protein, and a Ras protein or a nucleotide sequence capable of expressing Ras protein. In such methods, where the desired modulation is an inhibition of angiogenesis, at least one or both of the Raf or Ras proteins is inactive. Where the desired modulation is a stimulation of
20 angiogenesis, at least one or both of the Raf or Ras proteins are active.

Brief Description of the Drawings

 In the drawings forming a portion of this disclosure:

 FIGs. 1A-1D illustrate that ecotrophically packaged retrovirus only infects murine cells. Ecotrophic packaging cells were transfected with a retroviral construct
25 encoding the b-Galactosidase (b-Gal) gene and the supernatant collected 24 hours later. Supernate containing the virus was placed on either murine-derived fibroblasts (FIG. 1A), murine-derived endothelial cells (FIG. 1B), human epithelial adenocarcinoma cells (FIG. 1C), or human melanoma cells (FIG. 1D) for 24 hours. b-Gal activity was visualized using standard methods.

FIG. 2 illustrates that bFGF-induced increases in Raf activity were blocked by prior infection with Raf K375M in a mouse endothelial cell line. Ecotrophic packaging cells were transfected with a retroviral construct encoding the defective Raf kinase gene and the supernatant collected 24 hours later. Supernate containing virus was placed on mouse endothelial cells for 24 hours. Cells were then treated with bFGF for 5 minutes and lysed. Raf kinase activity was quantified by the ability of immunoprecipitated Raf kinase to phosphorylate the MEK substrate with radioactively labeled ^{32}P . Reaction mixtures were fractionated by SDS PAGE and quantified using scanning densitometry.

FIGS. 3A-3B illustrate that mutant inactive Raf K375M blocks bFGF-induced angiogenesis in a murine subcutaneous angiogenesis model. Angiogenesis was induced by injecting 250 μl of ice-cold, growth factor-reduced matrigel containing 400 ng/ml bFGF, with or without retrovirus expressing packaging cells that express Raf K375M, subcutaneously in the mouse flank. Five days later endothelial-specific FITC-conjugated Bandeiraea Simplifica B5 lectin was injected via the tail vein and allowed to circulate and clear for 30 minutes. Angiogenesis was then quantitated by removing, extracting, and assaying the angiogenic tissue for fluorescent content (FIG. 3A). Neovascularization was confirmed by optical sectioning (FIG. 3B).

FIGS. 4A-4B illustrate that mutationally active Raf stimulates angiogenesis in a murine subcutaneous angiogenesis model. Angiogenesis was induced by injecting 250 μl of ice-cold, growth factor-reduced matrigel containing retrovirus expressing packaging cells which express GFP control or amino terminal deleted Raf kinase (Raf 306-648), subcutaneously in the mouse flank. Five days later angiogenesis was then quantitated by removing, extracting, and assaying the angiogenic tissue for fluorescent content (FIG. 4A). Neovascularization was confirmed by sectioning and staining with Mason's trichrome (FIG. 4B).

FIGS. 5A-5D illustrate retroviral delivery of Raf K375M kinase to the tumor induced apoptosis in an endothelial-specific manner. Human tumors were injected subcutaneously on the flank of athymic wehi (nu/nu) mice and allowed to implant.

When tumors reached 100 mm³ they were injected intratumorally with culture supernate containing 10⁶ pfu of ecotrophically packaged Raf K375M. Forty-eight hours later the tumor was harvested, sectioned, and immunohistochemistry performed. Endothelial cells were identified by vWF expression (FIG. 5A), while the Flag tag marker was used to indicate cells infected by the Raf K375M kinase gene (FIG. 5B). Each of these markers are seen colocalized with the TUNEL marker indicative of apoptotic cells (FIGs. 5C & 5D).

FIGs. 6A-6B illustrate endothelial delivery of the Raf K375M kinase gene inhibited tumor growth and stimulated tumor regression. Human tumors were injected subcutaneously on the flank of athymic wehi (nu/nu) mice and allowed to grow to 100 mm³. At this point either a single injection of packaging cells expressing Raf K375M kinase was performed at a tumor-adjacent site or a series of intratumoral injections of viral supernate was initiated. This strategy resulted in rapid regressions of the tumors which was not seen with injection of the control GFP gene (FIG. 6A). This regression occurred rapidly and was maintained throughout the length of the experiment (FIG. 6B).

FIG. 7 depicts a cDNA sequence encoding for human c-Raf which is the complete coding sequence with the introns deleted. The sequence is accessible through GenBank Accession Number X03484 (GI=35841, HSRAFR). (SEQ ID NO.: 1).

FIG. 8 depicts the encoded translated amino acid residue sequence of human c-Raf of the coding sequence depicted in the nucleic acid sequence shown in FIG. 7. (SEQ ID NO.: 2).

FIG. 9 illustrates that angiogenesis is dependent on activation of the Ras-Raf-MEK-ERK pathway. Ras activity was elevated in chick chorioallantoic membrane (CAM) lysates exposed to bFGF as determined by a Ras pulldown assay. CAMs from 10-day old chick embryos were stimulated topically with filter disks saturated with either PBS or 30 nanograms (ng) of bFGF. After 5 minutes, CAM tissue was resected, homogenized in lysis buffer, and Ras activity was then determined

by its capacity to be precipitated by a GST fusion peptide encoding the Ras binding domain of Raf. Because only active Ras binds Raf, a recombinant protein was generated consisting of the Ras binding domain of Raf conjugated to glutathione-S-transferase (GST). In turn GST was conjugated to sepharose beads enabling the precipitation of active Ras from a tissue lysate.

FIG. 10 depicts the cDNA coding domain nucleotide sequence of wild-type human Ras (wt H-Ras). (SEQ ID NO.: 3). A complete coding sequence for c-Ha-Ras1 proto-oncogene is accessible through GenBank (GI=190890, HUMRASH). (SEQ ID NO.: 5).

FIG. 11 depicts the amino acid residue sequence encoded by the cDNA nucleotide sequence of wild-type human Ras (wt H-Ras) shown in FIG. 10. (SEQ ID NO.: 4).

FIG. 12 illustrates that infection with mutant null Ras blocked growth factor-induced angiogenesis in the CAM. Fifteen microliters (ul) of high titer Chicken sarcoma retrovirus, RCAS(A), encoding mutant null Ras, S17N Ras (wild type H-Ras with a substitution of Asn for Ser at position 17), was topically applied to filter disks on CAMs as stimulated with bFGF as described in FIG. 9. Angiogenesis was assessed after 72 hours by counting vessel branch points.

FIGs. 13A and 13B illustrate schematically and graphically respectively that infection with a mutant Ras construct, Ras V12S35, which selectively activates the Ras-Raf-MEK-ERK pathway, induced angiogenesis, whereas a mutant construct, Ras V12C40, which selectively activates the PI3K pathways, did not. Fifteen ul of high titer RCAS (A) virus encoding the Raf-MEK-ERK activating Ras construct, Ras V12S35, or the PI3 kinase activating Ras construct, Ras V12C40, were topically applied to filter disks and results assessed as described in FIG. 12.

FIG. 14 depicts the nucleotide sequence encoding the fusion protein Raf-caax, where the nucleotide sequence encoding the carboxy terminus of human Raf (wt H-Raf) is fused with a nucleotide sequence of encoding a 20 amino acid residue sequence of the K-Ras membrane localization domain. (SEQ ID NO.: 6).

FIG. 15 depicts the amino acid residue sequence of Raf-caax, the fusion protein generated from the fusion nucleotide sequence depicted in FIG. 14. (SEQ ID NO.: 7).

FIGs. 16A-16E and FIG. 16F, respectively, pictorially and graphically illustrate that the MEK inhibitor, PD98059, blocked angiogenesis induced by either mutant active Ras or Raf. Virus encoding the activating Ras construct, Ras V12 (also referred to as G12V, and the activating Raf construct, Raf-caax, were topically applied to filter disks as described in FIG. 12. After 24 hours, one (1) nanomole of the MEK inhibitor, PD98059, was added to the disk. The CAMs were then evaluated as described in FIG. 12. Data plotted is the mean \pm SE of 20 embryos.

FIGs. 17A-17F and FIG. 17G, respectively, pictorially and graphically illustrate that angiogenesis induced by Raf, but not Ras, was refractory to inhibition by integrin blockade. Infection with both mutant active Ras and Raf constructs induced pronounced angiogenesis, but only Ras-induced angiogenesis was inhibited by $\alpha_v\beta_3$ integrin-blocking antibodies. CAMs from 10-day old chick embryos were stimulated as described in FIGs. 9 and 12 with filter disks saturated with either PBS (control), bFGF, the RCAS(A) retroviral constructs G12V-Ras or Raf-caax. LM609, a monoclonal antibody to integrin $\alpha_v\beta_3$, was intravenously delivered after 24 hours and angiogenesis was assessed by vessel branch point analysis after 72 hours. Representative CAMs are shown in the inset. Data is the mean \pm SE of 20 embryos.

FIGs. 18A-18D and 18E, respectively, pictorially and graphically illustrate that co-infection of CAMs with a mutant null focal adhesion kinase, FRNK, blocked Ras, but not Raf-induced angiogenesis. RCAS(A) viruses encoding Ras V12 or Raf-caax were topically applied as described in FIG. 12 along with RCAS(B) virus encoding FAK-related-null-kinase (FRNK) to the CAM filter disk. Data is the mean \pm SE of 20 embryos.

FIGs. 19A and 19B-19G, respectively, graphically and pictorially, illustrate that FRNK blocked bFGF and Ras-, but not Raf, -induced angiogenesis in a murine subcutaneous angiogenesis model. Angiogenesis was induced by injecting 250 μ l of

ice-cold, growth factor-reduced matrigel containing either 400 ng/ml bFGF or Moloney retrovirus expressing packaging cells expressing the described gene, subcutaneously in the mouse flank. FRNK retrovirus was added to matrigel as high titer virus packaged with the vsv.g coat protein. Five days later, endothelial-specific FITC-conjugated Bandeiraea Simplicifolia B5 lectin was injected via the tail vein and allowed to circulate. Angiogenesis was then quantitated by removing, extracting, and assaying the angiogenic tissue for fluorescent content.

FIGs. 20A and 20B illustrate that co-infection of CAMs with a mutant null focal adhesion kinase, FRNK, blocked Ras-induced activation of Raf. CAMs were treated as described in FIG. 18 with the exception that after 24 hours the angiogenic tissue was resected, solubilized, Raf immunoprecipitated, and Raf activity assessed by its capacity to phosphorylate kinase-dead MEK. FIG. 20A shows the immunoprecipitated active versus total Raf proteins assayed under each of the combinations above the results. FIG. 20B graphically plots the results of the active Raf determinations under those conditions.

Detailed Description of the Invention

A. Definitions

Amino Acid Residue: An amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. The amino acid residues described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property is retained by the polypeptide. NH₂ refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature (described in J. Biol. Chem., 243:3552-59 (1969) and adopted at 37 CFR §1.822(b)(2)).

It should be noted that all amino acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the

beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues.

5 Polypeptide: refers to a linear series of amino acid residues connected to one another by peptide bonds between the alpha-amino group and carboxy group of contiguous amino acid residues.

Peptide: as used herein refers to a linear series of no more than about 50 amino acid residues connected one to the other as in a polypeptide.

10 Cyclic peptide: refers to a compound having a heteroatom ring structure that includes several amide bonds as in a typical peptide. The cyclic peptide can be a "head to tail" cyclized linear polypeptide in which a linear peptide's n-terminus has formed an amide bond with the -terminal carboxylate of the linear peptide, or it can contain a ring structure in which the polymer is homodetic or heterodetic and comprises amide bonds and/or other bonds to close the ring, such as disulfide bridges, thioesters, thioamides, guanidino, and the like linkages.

15 Protein: refers to a linear series of greater than 50 amino acid residues connected one to the other as in a polypeptide.

Fusion protein: refers to a polypeptide containing at least two different polypeptide domains operatively linked by a typical peptide bond ("fused"), where the two domains correspond to peptides not found fused in nature.

20 Synthetic peptide: refers to a chemically produced chain of amino acid residues linked together by peptide bonds that is free of naturally occurring proteins and fragments thereof.

B. General Considerations

25 The present invention relates generally to the discovery that angiogenesis is mediated by the protein kinase Raf protein, and that angiogenesis can be modulated by providing either active or inactive Raf proteins for potentiating or inhibiting angiogenesis, respectively. The invention also relates to the discovery that a Ras protein can affect Raf, and thereby modulate angiogenesis.

This discovery is important because of the role that angiogenesis, the formation of new blood vessels, plays in a variety of disease processes. On the other hand, where tissues associated with a disease condition require angiogenesis for tissue growth, it is desirable to inhibit angiogenesis and thereby inhibit the diseased tissue growth.

5 Where injured tissue requires angiogenesis for tissue growth and healing, it is desirable to potentiate or promote angiogenesis and thereby promote tissue healing and growth.

10 Where the growth of new blood vessels is the cause of, or contributes to, the pathology associated with a disease tissue, inhibition of angiogenesis will reduce the deleterious effects of the disease. By inhibiting angiogenesis, one can intervene in the disease, ameliorate the symptoms, and in some cases cure the disease.

15 Examples of tissue associated with disease and neovascularization that will benefit from inhibitory modulation of angiogenesis include cancer, rheumatoid arthritis, ocular diseases such as diabetic retinopathy, inflammatory diseases, restenosis, and the like. Where the growth of new blood vessels is required to support growth of a deleterious tissue, inhibition of angiogenesis reduces the blood supply to the tissue and thereby contributes to reduction in tissue mass based on blood supply requirements. Particularly preferred examples include growth of tumors where neovascularization is a continual requirement in order that the tumor grow beyond a
20 few millimeters in thickness, and for the establishment of solid tumor metastases.

25 Where the growth of new blood vessels contributes to healing of tissue, potentiation of angiogenesis assists in healing. Examples include treatment of patients with ischemic limbs in which there is abnormal, i.e. poor circulation as a result of diabetes or other conditions. Also contemplated are patients with chronic wounds which do not heal and therefore could benefit from the increase in vascular cell proliferation and neovascularization.

The methods of the present invention are effective in part because the therapy is highly selective for angiogenesis and not other biological processes.

As described earlier, angiogenesis includes a variety of processes involving neovascularization of a tissue including "sprouting", vasculogenesis, or vessel enlargement, all of which angiogenesis processes are affected by Raf protein alone or together with a Ras protein. With the exception of traumatic wound healing, corpus luteum formation and embryogenesis, it is believed that the majority of angiogenesis processes are associated with disease processes and therefore the use of the present therapeutic methods are selective for the disease and do not have deleterious side effects.

C. Raf Proteins

A protein kinase Raf protein for use in the present invention can vary depending upon the intended use. The terms "Raf protein" or "Raf" are used to refer collectively to the various forms of protein kinase Raf protein, either in active or inactive forms.

An "active Raf protein" refers to any of a variety of forms of Raf protein which potentiate, stimulate, activate, induce or increase angiogenesis. Assays to measure potentiation of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered active if the level of angiogenesis is at least 10% greater, preferably 25% greater, and more preferably 50% greater than a control level where no Raf is added to the assay system. The preferred assay for measuring potentiation is the *in vitro* Raf kinase as described in the Examples in which MEK substrate is phosphorylated with ^{32}P . Exemplary active Raf proteins are described in the Examples.

An "inactive Raf protein" refers to any of a variety of forms of Raf protein which inhibit, reduce, impede, or restrict angiogenesis. Assays to measure inhibition of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered inactive if the level of angiogenesis is at least 10% lower, preferably 25% lower, and more preferably 50% lower than a control level where no exogenous Raf is added to the assay system. The preferred assay for measuring

inhibition is the *in vitro* Raf kinase as described in the Examples in which MEK substrate is phosphorylated with ^{32}P . Exemplary inactive Raf proteins are described in the Examples.

5 A Raf protein useful in the present invention can be produced in any of a variety of methods including isolation from natural sources including tissue, production by recombinant DNA expression and purification, and the like. Raf protein can also be provided "in situ" by introduction of a gene therapy system to the tissue of interest which then expresses the protein in the tissue.

10 A gene encoding a Raf protein can be prepared by a variety of methods known in the art, and the invention is not to be construed as limiting in this regard. For example, the natural history of Raf is well known to include a variety of homologs from mammalian, avian, viral and the like species, and the gene can readily be cloned using cDNA cloning methods from any tissue expressing the protein. A preferred Raf for use in the invention is a cellular protein, such as the mammalian or avian homologs
15 designated c-Raf. Particularly preferred is human c-Raf. A further preferred Raf protein of this invention is a fusion protein of Raf that is constitutively active but independent of Ras-mediated activation. Such a Raf protein can be a fusion protein. A preferred Ras-independent Raf protein is Raf-caax which is a carboxy terminal fusion protein of wild type Raf with the K-Ras membrane localization domain as
20 further described in the Examples.

D. Ras Proteins

Ras family GTPases for use in the present invention can vary depending upon the intended use. The terms "Ras protein" or "Ras" are used herein to refer collectively to the various forms of Ras protein, either in active or inactive forms.

25 An "active Ras protein" refers to any of a variety of forms of Ras protein which potentiate, stimulate, activate, induce or increase angiogenesis. Assays to measure potentiation of angiogenesis by Ras are described herein, and are not to be construed as limiting. A protein is considered active if the level of angiogenesis is at

least 10% greater, preferably 25% greater, and more preferably 50% greater than a control level where no Ras is added to the assay system. Exemplary active Ras proteins are Ras G12V, also referred to as V12, and Ras V12S35, both of which are further described in the Examples.

5 An “inactive Ras protein” refers to any of a variety of forms of Ras protein which inhibit, impede, delay, or stop angiogenesis. Assays to measure inhibition of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered inactive if the level of angiogenesis is at least 10% lower, preferably 25% lower, and more preferably 50% lower than a control level where no exogenous Ras is added to the assay system. Exemplary inactive Ras proteins include the null mutant Ras referred to as Ras S17N (or sometimes N17) and V12C40, both of which are further described in the Examples.

10 A Ras protein useful in the present invention can be produced in any of a variety of methods including isolation from natural sources including tissue, production by recombinant DNA expression and purification, and the like. Ras protein can also be provided “in situ” by introduction of a gene therapy system to the tissue of interest which then expresses the protein in the tissue.

15 A gene encoding a Ras protein can be prepared by a variety of methods known in the art. The present invention is not to be construed as limiting in this regard. For example, the natural history of Ras is well known to include a variety of homologs from mammalian, avian, viral and the like species, and the gene can readily be cloned using cDNA cloning methods from any tissue expressing the protein.

20 It is to be understood by the present teachings that a Ras protein in its collective forms can be used in the same various embodiments as is described herein for a Raf protein, and therefore, the details for using a Ras protein are not reiterated. For example, Ras may be presented in an active or inactive form for modulating angiogenesis, or may be provided by nucleic acid expression of the Ras protein product, through the use of vector delivery systems, and in various pharmaceutical (therapeutic) compositions and articles of manufacture for practicing the invention.

0963700-0340

Methods of modulating angiogenesis using a Ras-based reagent in place of the recited Raf-based reagents are also contemplated.

E. Recombinant DNA Molecules and Expression Systems for Expression of a Raf or Ras Protein

5 The invention describes several nucleotide sequences of particular use in the present invention. These define nucleic acid sequences which encode for Raf or Ras protein useful in the invention, and various DNA segments, recombinant DNA (rDNA) molecules and vectors constructed for expression of Raf and/or Ras protein.

10 DNA molecules (segments) of this invention therefore can comprise sequences which encode whole structural genes, fragments of structural genes, and transcription units as described further herein.

 A preferred DNA segment is a nucleotide sequence which encodes a Raf protein as defined herein, or biologically active fragment thereof.

15 Another preferred DNA segment is a nucleotide sequence which encodes a Ras protein as defined herein, or biologically active fragment thereof. By biologically active, it is meant that the expressed protein will have at least some of the biological activity of the intact protein found in a cell, such as ligand binding, or in the case of active forms of the protein, enzymatic activity.

20 The amino acid residue sequence and nucleotide sequence of a preferred c-Raf and h-Ras are described in the Examples.

25 A preferred DNA segment codes for an amino acid residue sequence substantially the same as, and preferably consisting essentially of, an amino acid residue sequence or portions thereof corresponding to a Raf or Ras protein described herein. Representative and preferred DNA segments are further described in the Examples.

 The amino acid residue sequence of a protein or polypeptide is directly related via the genetic code to the deoxyribonucleic acid (DNA) sequence of the structural gene that codes for the protein. Thus, a structural gene or DNA segment can be

defined in terms of the amino acid residue sequence, i.e., protein or polypeptide, for which it codes.

An important and well known feature of the genetic code is its redundancy.

That is, for most of the amino acids used to make proteins, more than one coding

nucleotide triplet (codon) can code for or designate a particular amino acid residue.

Therefore, a number of different nucleotide sequences may code for a particular amino acid residue sequence. Such nucleotide sequences are functionally equivalent since they can result in the production of the same amino acid residue sequence in all

organisms. Occasionally, a methylated variant of a purine or pyrimidine may be incorporated into a given nucleotide sequence. However, such methylations do not affect the coding relationship in any way.

A nucleic acid is any polynucleotide or nucleic acid fragment, whether it be a polyribonucleotide or polydeoxyribonucleotide, i.e., RNA or DNA, or analogs thereof.

In preferred embodiments, a nucleic acid molecule is in the form of a segment of

duplex DNA, i.e., a DNA segment, although for certain molecular biological methodologies, single-stranded DNA or RNA is preferred.

DNA segments are produced by a number of means including chemical synthesis methods and recombinant approaches, preferably by cloning or by polymerase chain reaction (PCR). DNA segments that encode all or only portions of a

Raf or Ras protein can easily be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci et al, J. Am. Chem. Soc., 103:3185-3191 (1981), or using automated synthesis methods. In addition, larger DNA segments can readily be prepared by well known methods, such as synthesis of a group of oligonucleotides that define the DNA segment, followed by hybridization and ligation of

oligonucleotides to build the complete segment. Alternative methods include isolation of a preferred DNA segment by PCR with a pair of oligonucleotide primers used on a cDNA library believed to contain members which encode a Raf or Ras protein.

Of course, through chemical synthesis, any desired modifications can be made simply by substituting the appropriate bases for those encoding the native amino acid

residue sequence. This method is well known, and can be readily applied to the production of the various different "modified" Raf or Ras proteins described herein.

Furthermore, DNA segments consisting essentially of structural genes encoding a Raf or Ras protein can be subsequently modified, as by site-directed or random mutagenesis, to introduce any desired substitutions. It is understood that various allelic forms of Raf or Ras protein and genes encoding for Raf or Ras protein are also suitable for use in the present invention.

1. Cloning a Raf or Ras Gene

A Raf or Ras gene of this invention can be cloned from a suitable source of genomic DNA or messenger RNA (mRNA) by a variety of biochemical methods. Cloning these genes can be conducted according to the general methods described in the Examples and as known in the art.

Sources of nucleic acids for cloning a Raf or Ras gene suitable for use in the methods of this invention can include genomic DNA or messenger RNA (mRNA) in the form of a cDNA library, from a tissue believed to express these proteins. A preferred tissue is human lung tissue, although any other suitable tissue may be used.

A preferred cloning method involves the preparation of a cDNA library using standard methods, and isolating the Raf-encoding or Ras-encoding nucleotide sequence by PCR amplification using paired oligonucleotide primers based on the nucleotide sequences described herein. Alternatively, the desired cDNA clones can be identified and isolated from a cDNA or genomic library by conventional nucleic acid hybridization methods using a hybridization probe based on the nucleic acid sequences described herein. Other methods of isolating and cloning suitable Raf-encoding or Ras-encoding nucleic acids are readily apparent to one skilled in the art.

2. Expression Vectors

The invention contemplates a recombinant DNA molecule (rDNA) containing a DNA segment encoding a Raf and/or Ras protein as described herein. An

expressible rDNA can be produced by operatively (in frame, expressibly) linking a vector to a Raf or Ras encoding DNA segment of the present invention. It is envisioned that a combination expression can be constructed wherein Raf encoding and Ras encoding nucleic acid are present, either operably linked to the same, or
5 separate promoters. Thus, a recombinant DNA molecule is a hybrid DNA molecule comprising at least two nucleic acids of a nucleotide sequences not normally found together in nature (i.e. gene and vector).

The choice of vector to which a DNA segment of the present invention is operatively linked depends directly, as is well known in the art, on the functional
10 properties desired, e.g., protein expression, and the host cell to be transformed. Typical considerations in the art of constructing recombinant DNA molecules. A vector contemplated by the present invention is at least capable of directing the replication, and preferably also expression, of a structural gene included in the vector DNA segments, to which it is operatively linked.

Both prokaryotic and eukaryotic expression vectors are familiar to one of
15 ordinary skill in the art of vector construction, and are described by Ausubel, et al., in Current Protocols in Molecular Biology, Wiley and Sons, New York (1993) and by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory (1989). These references also describe many of the general recombinant
20 DNA methods referred to herein.

In one embodiment, a vector contemplated by the present invention includes a procaryotic replicon, i.e., a DNA sequence having the ability to direct autonomous replication and maintenance of the recombinant DNA molecule extrachromosomally in a procaryotic host cell, such as a bacterial host cell, transformed therewith. Such
25 replicons are well known in the art. In addition, those embodiments that include a procaryotic replicon also include a gene whose expression confers drug resistance to a bacterial host transformed therewith. Typical bacterial drug resistance genes are those that confer resistance to ampicillin or tetracycline.

Those vectors that include a procaryotic replicon can also include a procaryotic promoter capable of directing the expression (transcription and translation) of a structural gene in a bacterial host cell, such as E. coli, transformed therewith. A promoter is an expression control element formed by a DNA sequence that permits binding of RNA polymerase and transcription to occur. Promoters or other such regulatory nucleic acid sequences can be inducible or constitutive depending upon the desired expression control and/or effect. Promoter sequences compatible with bacterial hosts are typically provided in plasmid vectors containing convenient restriction sites for insertion of a DNA segment of the present invention. Typical of such vector plasmids are pUC8, pUC9, pBR322 and pBR329 available from Biorad Laboratories, (Richmond, CA), pRSET available from Invitrogen (San Diego, CA) and pPL and pKK223 available from Pharmacia, Piscataway, N.J.

Expression vectors compatible with eukaryotic cells, preferably those compatible with vertebrate cells, can also be used to form the recombinant DNA molecules of the present invention. Eukaryotic cell expression vectors are well known in the art and are available from several commercial sources. Typically, such vectors are provided containing convenient restriction sites for insertion of the desired DNA segment. Typical of such vectors are pSVL and pKSV-10 (Pharmacia), pBPV-1/pML2d (International Biotechnologies, Inc.), pTDT1 (ATCC, #31255), pRc/CMV (Invitrogen, Inc.), the preferred vector described in the Examples, and the like eukaryotic expression vectors.

A particularly preferred system for gene expression in the context of this invention includes a gene delivery component, that is, the ability to deliver the gene to the tissue of interest. Suitable vectors are "infectious" vectors such as recombinant DNA viruses, adenovirus or retrovirus vectors which are engineered to express the desired protein and have features which allow infection of preselected target tissues. Particularly preferred is the retrovirus vector system described herein.

Mammalian cell systems that utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus expression

vectors, the coding sequence of a polypeptide may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted into the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the polypeptide in infected hosts (e.g., see Logan et al., Proc. Natl. Acad. Sci., USA, 81:3655-3659 (1984)). Alternatively, the vaccinia virus 7.5K promoter may be used (e.g., see, Mackett et al., Proc. Natl. Acad. Sci., USA, 79:7415-7419 (1982); Mackett et al., J. Virol., 49:857-864 (1984); Panicali et al., Proc. Natl. Acad. Sci., USA, 79:4927-4931 (1982)). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver et al., Mol. Cell. Biol., 1:486 (1981)). Shortly after entry of this DNA into target cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by including a selectable marker in the plasmid, such as the neo gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the polypeptide-encoding nucleotide sequence in host cells (Cone et al., Proc. Natl. Acad. Sci., USA, 81:6349-6353 (1984)). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothionine IIA promoter and heat shock promoters.

Recently, long-term survival of cytomegalovirus (CMV) promoter versus Rous sarcoma virus (RSV) promotor-driven thymidine kinase (TK) gene therapy in nude mice bearing human ovarian cancer has been studied. Cell killing efficacy of adenovirus-mediated CMV promoter-driven herpes simplex virus TK gene therapy was found to be 2 to 10 times more effective than RSV driven therapy (Tong et al., Hybridoma 18(1):93-97 (1999)). The design of chimeric promoters for gene therapy applications, which call for low level expression followed by inducible high-level

expression has also been described (Suzuki et al., Human Gene Therapy 7:1883-1893 (1996)).

For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with a cDNA controlled by appropriate expression control elements (e.g., promoter and enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. As mentioned above, the selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell, 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska et al, Proc. Natl. Acad. Sci., USA, 48:2026 (1962)), and adenine phosphoribosyltransferase (Lowy et al., Cell, 22:817 (1980)) genes, which can be employed in tk⁻, hgp^rt⁻ or apr^t⁻ cells respectively. Also, antimetabolite resistance-conferring genes can be used as the basis of selection; for example, the genes for dhfr, which confers resistance to methotrexate (Wigler et al., Proc. Natl. Acad. Sci., USA, 77:3567 (1980); O'Hare et al., Proc. Natl. Acad. Sci., USA, 78:1527 (1981); gpt, which confers resistance to mycophenolic acid (Mulligan et al, Proc. Natl. Acad. Sci., USA, 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al, J. Mol. Biol., 150:1 (1981)); and hyg^r, which confers resistance to hygromycin (Santerre et al, Gene, 30:147 (1984)). Recently, additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman et al, Proc. Natl. Acad. Sci., USA, 85:804 (1988)); and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO

(McConlogue L., In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed. (1987)).

5 The principal vectors contemplated for human gene therapy, are derived from retroviral origin (Wilson, Clin. Exp. Immunol. 107(Sup. 1):31-32 (1997); Bank et al., Bioessays 18(12):999-1007 (1996); Robbins et al., Pharmacol. Ther. 80(1):35-47 (1998)). The therapeutic potential of gene transfer and antisense therapy has stimulated the development of many vector systems for treating a variety of tissues (vasculature, Stephan et al., Fundam. Clin. Pharmacol. 11(2):97-110 (1997); Feldman et al., Cardiovasc. Res. 35(3):391-404 (1997); Vassalli et al., Cardiovasc. Res. 10 35(3):459-69 (1997); Baek et al., Circ. Res. 82(3):295-305 (1998); kidney, Lien et al., Kidney Int. Suppl. 61:S85-8 (1997); liver, Ferry et al., Hum Gene Ther. 9(14):1975-81 (1998); muscle, Marshall et al., Curr. Opin. Genet. Dev. 8(3):360-5 (1998)). In addition to these tissues, a critical target for human gene therapy is cancer, either the tumor itself, or associated tissues. (Runnebaum, Anticancer Res. 17(4B):2887-90 15 (1997); Spear et al., J. Neurovirol. 4(2):133-47 (1998)).

Specific examples of viral gene therapy vector systems readily adaptable for use in the methods of the present invention are briefly described below. Retroviral gene delivery has been recently reviewed by Federspiel and Hughes (Methods in Cell Biol. 52:179-214 (1998)) which describes in particular, the avian leukosis virus (ALV) 20 retrovirus family (Federspiel et al., Proc. Natl. Acad. Sci., USA, 93:4931 (1996); Federspiel et al., Proc. Natl. Acad. Sci., USA, 91:11241 (1994)). Retroviral vectors, including ALV and murine leukemia virus (MLV) are further described by Svoboda (Gene 206:153-163 (1998)).

25 Modified retroviral/adenoviral expression systems can be readily adapted for practice of the methods of the present invention. For example, murine leukemia virus (MLV) systems are reviewed by Karavanas et al., Crit. Rev. in Oncology/Hematology, 28:7-30 (1998). Adenovirus expression systems are reviewed by Von Seggern and Nemerow in Gene Expression Systems (ed. Fernandez & Hoeffler, Academic Press, San Diego, CA, chapter 5, pages 112-157 (1999)).

Protein expression systems have been demonstrated to have effective use both *in vivo* and *in vitro*. For example, efficient gene transfer to human squamous cell carcinomas by a herpes simplex virus (HSV) type 1 amplicon vector has been described. (Carew et al., 1998, Am. J. Surg. 176:404-408). Herpes simplex virus has been used for gene transfer to the nervous system (Goins et al., J. Neurovirol. 3 (Sup. 1):S80-8 (1997)). Targeted suicide vectors using HSV-TK has been tested on solid tumors (Smiley et al., Hum. Gene Ther. 8(8):965-77 (1997)). Herpes simplex virus type 1 vector has been used for cancer gene therapy on colon carcinoma cells (Yoon et al., Ann. Surg. 228(3):366-74 (1998)). Hybrid vectors have been developed to extend the length of time of transfection, including HSV/AAV (adeno-associated virus) hybrids for treating hepatocytes (Fraefel et al., Mol. Med. 3(12):813-825 (1997)).

Vaccinia virus has been developed for human gene therapy because of its large genome (Peplinski et al., Surg. Oncol. Clin. N. Am. 7(3):575-88 (1998)). Thymidine kinase-deleted vaccinia virus expressing purine nucleoside pyrophosphorylase has been described for use as a tumor directed gene therapy vector. (Puhlman et al., Human Gene Therapy 10:649-657 (1999)).

Adeno-associated virus 2 (AAV) has been described for use in human gene therapy, however AAV requires a helper virus (such as adenovirus or herpes virus) for optimal replication and packaging in mammalian cells (Snoeck et al., Exp. Nephrol. 5(6):514-20 (1997); Rabinowitz et al., Curr. Opin. Biotechnol. 9(5):470-5 (1998)). However, *in vitro* packaging of an infectious recombinant AAV has been described, making this system much more promising (Ding et al., Gene Therapy 4:1167-1172 (1997)). It has been shown that the AAV mediated transfer of ecotropic retrovirus receptor cDNA allows ecotropic retroviral transduction of established and primary human cells (Qing et al., J. Virology 71(7):5663-5667 (1997)). Cancer gene therapy using an AAV vector expressing human wild-type p53 has been demonstrated (Qazilbash et al., Gene Therapy 4:675-682 (1997)). Gene transfer into vascular cells using AAV vectors has also been shown (Maeda et al., Cardiovascular Res. 35:514-521 (1997)). AAV has been demonstrated as a suitable vector for liver directed gene

therapy (Xiao et al., J. Virol. 72(12):10222-6 (1998)). AAV vectors have been demonstrated for use in gene therapy of brain tissues and the central nervous system (Chamberlin et al., Brain Res. 793(1-2):169-75 (1998); During et al., Gene Therapy 5(6):820-7 (1998)). AAV vectors have also been compared with adenovirus vectors (AdV) for gene therapy of the lung and transfer to human cystic fibrosis epithelial cells (Teramoto et al., J. Virol. 72(11):8904-12 (1998)).

Chimeric AdV/retroviral gene therapy vector systems which incorporate the useful qualities of each virus to create a nonintegrative AdV that is rendered functionally integrative via the intermediate generation of a retroviral producer cell (Feng et al., Nat. Biotechnology 15(9):866-70 (1997); Bilbao et al., FASEB J 11(8):624-34 (1997)). This powerful new generation of gene therapy vector has been adapted for targeted cancer gene therapy (Bilbao et al., Adv. Exp. Med. Biol. 451:365-74 (1998)). Single injection of AdV expressing p53 inhibited growth of subcutaneous tumor nodules of human prostate cancer cells (Asgari et al., Int. J. Cancer 71(3):377-82 (1997)). AdV mediated gene transfer of wild-type p53 in patients with advanced non-small cell lung cancer has been described (Schuler et al., Human Gene Therapy 9:2075-2082 (1998)). This same cancer has been the subject of p53 gene replacement therapy mediated by AdV vectors (Roth et al., Semin. Oncol. 25(3 Suppl 8):33-7 (1998)). AdV mediated gene transfer of p53 inhibits endothelial cell differentiation and angiogenesis *in vivo* (Riccioni et al., Gene Ther. 5(6):747-54 (1998)). Adenovirus-mediated expression of melanoma antigen gp75 as immunotherapy for metastatic melanoma has also been described (Hirschowitz et al., Gene Therapy 5:975-983 (1998)). AdV facilitates infection of human cells with ecotropic retrovirus and increases efficiency of retroviral infection (Scott-Taylor, et al., Gene Ther. 5(5):621-9 (1998)). AdV vectors have been used for gene transfer to vascular smooth muscle cells (Li et al., Chin. Med. J.(Engl) 110(12):950-4 (1997)), squamous cell carcinoma cells (Goebel et al., Otolaryngol Head Neck Surg 119(4):331-6 (1998)), esophageal cancer cells (Senmaru et al., Int J. Cancer 78(3):366-71 (1998)), mesangial cells (Nahman et al., J. Investig. Med. 46(5):204-9 (1998)), glial cells (Chen et al.,

Cancer Res. 58(16):3504-7 (1998)), and to the joints of animals (Ikeda et al., J. Rheumatol. 25(9):1666-73 (1998)). More recently, catheter-based pericardial gene transfer mediated by AcV vectors has been demonstrated (March et al., Clin. Cardiol. 22(1 Suppl 1):I23-9 (1999)). Manipulation of the AdV system with the proper
5 controlling genetic elements allows for the AdV-mediated regulatable target gene expression *in vivo* (Burcin et al., PNAS (USA) 96(2):355-60 (1999)).

Alphavirus vectors have been developed for human gene therapy applications, with packaging cell lines suitable for transformation with expression cassettes suitable for use with Sindbis virus and Semliki Forest virus-derived vectors (Polo et al., Proc. Natl. Acad. Sci., USA, 96:4598-4603 (1999)). Noncytopathic flavivirus replicon
10 RNA-based systems have also been developed (Varnavski et al., Virology 255(2):366-75 (1999)). Suicide HSV-TK gene containing sinbis virus vectors have been used for cell-specific targeting into tumor cells (Iijima et al., Int. J. Cancer 80(1):110-8 (1998)).

Retroviral vectors based on human foamy virus (HFV) also show promise as
15 gene therapy vectors (Trowbridge et al., Human Gene Therapy 9:2517-2525 (1998)). Foamy virus vectors have been designed for suicide gene therapy (Nestler et al., Gene Ther. 4(11):1270-7 (1997)). Recombinant murine cytomegalovirus and promoter systems have also been used as vectors for high level expression (Manning et al., J. Virol. Meth. 73(1):31-9 (1998); Tong et al., Hybridoma 18(1):93-7 (1998)).

20 Gene delivery into non-dividing cells has been made feasible by the generation of Sendai virus based vectors (Nakanishi et al., J. Controlled Release 54(1):61-8 (1998)).

In other efforts to enable the transformation of non-dividing somatic cells, lentiviral vectors have been explored. Gene therapy of cystic fibrosis using a
25 replication-defective human immunodeficiency virus (HIV) based vector has been described. (Goldman et al., Human Gene Therapy 8:2261-2268 (1997)). Sustained expression of genes delivered into liver and muscle by lentiviral vectors has also been shown (Kafri et al., Nat. Genet. 17(3):314-7 (1997)). However, safety concerns are predominant, and improved vector development is proceeding rapidly (Kim et al., J.

Virology 72(2):994-1004 (1998)). Examination of the HIV LTR and Tat yield important information about the organization of the genome for developing vectors (Sadaie et al., J. Med. Virol. 54(2):118-28 (1998)). Thus, the genetic requirements for an effective HIV based vector are now better understood (Gasmi et al., J. Virol. 73(3):1828-34 (1999)). Self inactivating vectors, or conditional packaging cell lines have been described (for example, Zuffery et al., J. Virol. 72(12):9873-80 (1998); Miyoshi et al., J. Virol. 72(10):8150-7 (1998); Dull et al., J. Virol. 72(11):8463-71 (1998); and Kaul et al., Virology 249(1):167-74 (1998)). Efficient transduction of human lymphocytes and CD34+ cells by HIV vectors has been shown (Douglas et al., Hum. Gene Ther. 10(6):935-45 (1999); Miyoshi et al., Science 283(5402):682-6 (1999)). Efficient transduction of nondividing human cells by feline immunodeficiency virus (FIV) lentiviral vectors has been described, which minimizes safety concerns with using HIV based vectors (Poeschla et al., Nature Medicine 4(3):354-357 (1998)). Productive infection of human blood mononuclear cells by FIV vectors has been shown (Johnston et al., J. Virol. 73(3):2491-8 (1999)).

While many viral vectors are difficult to handle, and capacity for inserted DNA limited, these limitations and disadvantages have been addressed. For example, in addition to simplified viral packaging cell lines, Mini-viral vectors, derived from human herpes virus, herpes simplex virus type 1 (HSV-1), and Epstein-Barr virus (EBV), have been developed to simplify manipulation of genetic material and generation of viral vectors (Wang et al., J. Virology 70(12):8422-8430 (1996)). Adaptor plasmids have been previously shown to simplify insertion of foreign DNA into helper-independent Retroviral vectors (J. Virology 61(10):3004-3012 (1987)).

Viral vectors are not the only means for effecting gene therapy, as several non-viral vectors have also been described. A targeted non-viral gene delivery vector based on the use of Epidermal Growth Factor/DNA polyplex (EGF/DNA) has been shown to result in efficient and specific gene delivery (Cristiano, Anticancer Res. 18:3241-3246 (1998)). Gene therapy of the vasculature and CNS have been demonstrated using cationic liposomes (Yang et al., J. Neurotrauma 14(5):281-97

(1997)). Transient gene therapy of pancreatitis has also been accomplished using cationic liposomes (Denham et al., Ann. Surg. 227(6):812-20 (1998)). A chitosan-based vector/DNA complexes for gene delivery have been shown to be effective (Erbacher et al., Pharm. Res. 15(9):1332-9 (1998)). A non-viral DNA delivery vector based on a terplex system has been described (Kim et al., 53(1-3):175-82 (1998)). Virus particle coated liposome complexes have also been used to effect gene transfer (Hirai et al., Biochem. Biophys. Res. Commun. 241(1):112-8 (1997)).

Cancer gene therapy by direct tumor injections of nonviral T7 vector encoding a thymidine kinase gene has been demonstrated (Chen et al., Human Gene Therapy 9:729-736 (1998)). Plasmid DNA preparation is important for direct injection gene transfer (Horn et al., Hum. Gene Ther. 6(5):656-73 (1995)). Modified plasmid vectors have been adapted specifically for direct injection (Hartikka et al., Hum. Gene Ther. 7(10):1205-17 (1996)).

Thus, a wide variety of gene transfer/gene therapy vectors and constructs are known in the art. These vectors are readily adapted for use in the methods of the present invention. By the appropriate manipulation using recombinant DNA/molecular biology techniques to insert an operatively linked Raf or Ras encoding nucleic acid segment (either active or inactive) into the selected expression/delivery vector, many equivalent vectors for the practice of the present invention can be generated.

F. Methods For Modulation of Angiogenesis

In one aspect, the present invention provides for a method for the modulation of angiogenesis in a tissue associated with a disease process or condition, and thereby affect events in the tissue which depend upon angiogenesis. Generally, the method comprises administering to the tissue, associated with, or suffering from a disease process or condition, an angiogenesis-modulating amount of a composition comprising a Raf protein or a nucleic acid vector expressing active or inactive Raf.

A further method comprises administering to the tissue, associated with a disease process or condition, an angiogenesis-modulating amount of a composition comprising a Ras protein or a nucleic acid vector expressing active or inactive Ras. Another method aspect comprises administering to the tissue associated with a disease process or condition, an angiogenesis-modulating amount of a Raf and Ras protein or one or more nucleic acid vector expressing active or inactive Raf and Ras.

Any of a variety of tissues, or organs comprised of organized tissues, can support angiogenesis in disease conditions including skin, muscle, gut, connective tissue, brain tissue, nerve cells, joints, bones and the like tissue in which blood vessels can invade upon angiogenic stimuli.

The patient to be treated according to the present invention in its many embodiments is a human patient, although the invention is effective with respect to all mammals. In this context, a "patient" is a human patient as well as a veterinary patient, a mammal of any mammalian species in which treatment of tissue associated with diseases involving angiogenesis is desirable, particularly agricultural and domestic mammalian species.

Thus, the method embodying the present invention comprises administering to a patient a therapeutically effective amount of a physiologically tolerable composition containing a Raf and/or Ras protein or nucleic acid vector for expressing a Raf and/or Ras protein.

The dosage ranges for the administration of a Raf or Ras protein depend upon the form of the protein, and its potency, as described further herein, and are amounts large enough to produce the desired effect in which angiogenesis and the disease symptoms mediated by angiogenesis are ameliorated. The dosage should not be so large as to cause adverse side effects, such as hyperviscosity syndromes, pulmonary edema, congestive heart failure, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient and can be determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication.

A therapeutically effective amount is an amount of Raf or Ras protein, or nucleic acid encoding for (active or inactive) Raf or Ras protein, sufficient to produce a measurable modulation of angiogenesis in the tissue being treated, i.e., an angiogenesis-modulating amount. Modulation of angiogenesis can be measured or monitored *in vitro* by CAM assay as described herein, examination of tumor tissues, or by other methods known to one skilled in the art.

The Raf or Ras protein or nucleic acid vector expressing such protein can be administered parenterally by injection or by gradual infusion over time. Although the tissue to be treated can typically be accessed in the body by systemic administration and therefore most often treated by intravenous administration of therapeutic compositions, other tissues and delivery means are contemplated where there is a likelihood that the tissue targeted contains the target molecule. Thus, compositions of the invention can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, transdermally, and can be delivered by peristaltic means, if desired.

The therapeutic compositions containing a Raf or Ras protein or nucleic acid vector expressing the Raf or Ras protein can be conventionally administered intravenously, as by injection of a unit dose, for example. The term "unit dose" when used in reference to a therapeutic composition of the present invention refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required physiologically acceptable diluent; i.e., carrier, or vehicle.

In one preferred embodiment the active material is administered in a single dosage intravenously. Localized administration can be accomplished by direct injection or by taking advantage of anatomically isolated compartments, isolating the microcirculation of target organ systems, reperfusion in a circulating system, or catheter based temporary occlusion of target regions of vasculature associated with diseased tissues.

The compositions are administered in a manner compatible with the dosage formulation, and in a therapeutically effective amount. The quantity to be administered and timing depends on the subject to be treated, capacity of the subject's system to utilize the active ingredient, and degree of therapeutic effect desired.

5 Precise amounts of active ingredient required to be administered depend on the judgement of the practitioner and are peculiar to each individual. However, suitable dosage ranges for systemic application are disclosed herein and depend on the route of administration. Suitable regimes for administration are also variable, but are typified by an initial administration followed by repeated doses at one or more hour intervals
10 by a subsequent injection or other administration. Alternatively, continuous intravenous infusion sufficient to maintain concentrations in the blood in the ranges specified for *in vivo* therapies are contemplated.

1. Inhibition of Angiogenesis

There are a variety of diseases in which inhibition of angiogenesis is
15 important, referred to as angiogenic diseases, including but not limited to, inflammatory disorders such as immune and non-immune inflammation, chronic articular rheumatism and psoriasis, disorders associated with inappropriate or inopportune invasion of vessels such as diabetic retinopathy, neovascular glaucoma, restenosis, capillary proliferation in atherosclerotic plaques and osteoporosis, and
20 cancer associated disorders, such as solid tumors, solid tumor metastases, angiofibromas, retrolental fibroplasia, hemangiomas, Kaposi sarcoma and the like cancers which require neovascularization to support tumor growth.

Thus, methods which inhibit angiogenesis in a tissue associated with a disease condition ameliorates symptoms of the disease and, depending upon the disease, can
25 contribute to cure of the disease. In one embodiment, the invention contemplates inhibition of angiogenesis, per se, in a tissue associated with a disease condition. The extent of angiogenesis in a tissue, and therefore the extent of inhibition achieved by the present methods, can be evaluated by a variety of methods.

Thus, in one embodiment, a tissue to be treated is an inflamed tissue and the angiogenesis to be inhibited is inflamed tissue angiogenesis where there is neovascularization of inflamed tissue. This particular method includes inhibition of angiogenesis in arthritic tissues, such as in a patient with chronic articular rheumatism, in immune or non-immune inflamed tissues, in psoriatic tissue, and the like.

In another embodiment, a tissue to be treated is a retinal tissue of a patient suffering from a retinal disease such as diabetic retinopathy, macular degeneration or neovascular glaucoma and the angiogenesis to be inhibited is retinal tissue angiogenesis where there is neovascularization of retinal tissue.

In an additional embodiment, a tissue to be treated is a tumor tissue of a patient with a solid tumor, a metastases, a skin cancer, a breast cancer, a hemangioma or angiofibroma and the like cancer, and the angiogenesis to be inhibited is tumor tissue angiogenesis where there is neovascularization of a tumor tissue. Typical solid tumor tissues treatable by the present methods include lung, pancreas, breast, colon, laryngeal, ovarian, and the like tissues. Inhibition of tumor tissue angiogenesis is a particularly preferred embodiment because of the important role neovascularization plays in tumor growth. In the absence of neovascularization of tumor tissue, the tumor tissue does not obtain the required nutrients, slows in growth, ceases additional growth, regresses and ultimately becomes necrotic resulting in killing of the tumor.

Stated in other words, the present invention provides for a method of inhibiting tumor neovascularization by inhibiting tumor angiogenesis according to the present methods. Similarly, the invention provides a method of inhibiting tumor growth by practicing the angiogenesis-inhibiting methods.

The methods are also particularly effective against the formation of metastases because (1) their formation requires vascularization of a primary tumor so that the metastatic cancer cells can exit the primary tumor and (2) their establishment in a secondary site requires neovascularization to support growth of the metastases.

In a yet further embodiment, the invention contemplates the practice of the method in conjunction with other therapies such as conventional chemotherapy

directed against solid tumors and for control of establishment of metastases. The administration of angiogenesis inhibitor is typically conducted during or after chemotherapy, although it is preferably to inhibit angiogenesis after a regimen of chemotherapy at times where the tumor tissue will be responding to the toxic assault by inducing angiogenesis to recover by the provision of a blood supply and nutrients to the tumor tissue. In addition, it is preferred to administer the angiogenesis inhibition methods after surgery where solid tumors have been removed as a prophylaxis against metastases.

Insofar as the present methods apply to inhibition of tumor neovascularization, the methods can also apply to inhibition of tumor tissue growth, to inhibition of tumor metastases formation, and to regression of established tumors.

Restenosis is a process of smooth muscle cell (SMC) migration and proliferation into the tissue at the site of percutaneous transluminal coronary angioplasty which hampers the success of angioplasty. The migration and proliferation of SMC's during restenosis can be considered a process of angiogenesis which is inhibited by the present methods. Therefore, the invention also contemplates inhibition of restenosis by inhibiting angiogenesis according to the present methods in a patient following angioplasty procedures. For inhibition of restenosis, the inactivated tyrosine kinase is typically administered after the angioplasty procedure because the coronary vessel wall is at risk of restenosis, typically for from about 2 to about 28 days, and more typically for about the first 14 days following the procedure.

The present method for inhibiting angiogenesis in a tissue associated with a disease condition, and therefore for also practicing the methods for treatment of angiogenesis-related diseases, comprises contacting a tissue in which angiogenesis is occurring, or is at risk for occurring, with a therapeutically effective amount of a composition comprising an inactivated Raf protein or vector expressing the protein. Inhibition of angiogenesis and tumor regression occurs as early as 7 days after the initial contacting with the therapeutic composition. Additional or prolonged exposure to inactive Raf or Ras protein is preferable for 7 days to 6 weeks, preferably about 14

to 28 days. Shorter periods of exposure can be useful where the modulating effects are detectable earlier, however administration and subsequent exposure for at least 12 hours is preferred.

2. Potentialiation of Angiogenesis

5 In cases where it is desirable to promote or potentiate angiogenesis, administration of an active Raf or Ras protein to the tissue is useful. The routes and timing of administration are comparable to the methods described hereinabove for inhibition.

G. Therapeutic Compositions

10 The present invention contemplates therapeutic compositions useful for practicing the therapeutic methods described herein. Therapeutic compositions of the present invention contain a physiologically tolerable carrier together with a Raf or Ras protein or vector capable of expressing a Raf or Ras protein as described herein, dissolved or dispersed therein as an active ingredient. In a preferred embodiment, the
15 therapeutic composition is not immunogenic when administered to a mammal or human patient for therapeutic purposes.

As used herein, the terms "pharmaceutically acceptable", "physiologically tolerable" and grammatical variations thereof, as they refer to compositions, carriers, diluents and reagents, are used interchangeably and represent that the materials are
20 capable of administration to or upon a mammal without the production of undesirable physiological effects such as nausea, dizziness, gastric upset and the like.

The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art and need not be limited based on formulation. Typically such compositions are prepared as injectable
25 either as liquid solutions or suspensions, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be emulsified or presented as a liposome composition.

The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if
5 desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like which enhance the effectiveness of the active ingredient.

The therapeutic composition of the present invention can include pharmaceutically acceptable salts of the components therein. Pharmaceutically
10 acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium or
15 ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

Physiologically tolerable carriers are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at
20 physiological pH value, physiological saline or both, such as phosphate-buffered saline. Still further, aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and potassium chlorides, dextrose, polyethylene glycol and other solutes.

Liquid compositions can also contain liquid phases in addition to and to the
25 exclusion of water. Exemplary of such additional liquid phases are glycerin, vegetable oils such as cottonseed oil, and water-oil emulsions.

A therapeutic composition contains an angiogenesis-modulating amount of a Raf or Ras protein of the present invention, or sufficient recombinant DNA expression vector to express an effective amount of Raf or Ras protein, typically formulated to

contain an amount of at least 0.1 weight percent of Raf or Ras protein per weight of total therapeutic composition. A weight percent is a ratio by weight of Raf protein to total composition. Thus, for example, 0.1 weight percent is 0.1 grams of Raf or Ras protein per 100 grams of total composition. For DNA expression vectors, the amount administered depends on the properties of the expression vector, the tissue to be treated, and the like considerations. The suitable amount administered can be measured by amount of vector, or amount of expressed protein that is expected.

H. Article of Manufacture

The invention also contemplates an article of manufacture which is a labeled container for providing a Raf or Ras protein of the invention. An article of manufacture comprises packaging material and a pharmaceutical agent contained within the packaging material.

The pharmaceutical agent in an article of manufacture is any of the compositions of the present invention suitable for providing a Raf or Ras protein and formulated into a pharmaceutically acceptable form as described herein according to the disclosed indications. Thus, the composition can comprise a Raf and/or Ras protein or a DNA molecule which is capable of expressing a Raf and/or Ras protein. The article of manufacture contains an amount of pharmaceutical agent sufficient for use in treating a condition indicated herein, either in unit or multiple dosages.

The packaging material comprises a label which indicates the use of the pharmaceutical agent contained therein, e.g., for treating conditions assisted by the inhibition or potentiation of angiogenesis, and the like conditions disclosed herein. The label can further include instructions for use and related information as may be required for marketing. The packaging material can include container(s) for storage of the pharmaceutical agent.

As used herein, the term packaging material refers to a material such as glass, plastic, paper, foil, and the like capable of holding within fixed means a pharmaceutical agent. Thus, for example, the packaging material can be plastic or

glass vials, laminated envelopes and the like containers used to contain a pharmaceutical composition including the pharmaceutical agent.

In preferred embodiments, the packaging material includes a label that is a tangible expression describing the contents of the article of manufacture and the use of the pharmaceutical agent contained therein.

Examples

The following examples relating to this invention are illustrative and should not, of course, be construed as specifically limiting the invention. Moreover, such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are to be considered to fall within the scope of the present invention hereinafter claimed.

1. Preparation of c-Raf Expression Constructs

For preparing the expression constructs useful in modulating angiogenesis by the methods of the present invention, c-Raf cDNA is manipulated and inserted into an expression construct/vector.

The cDNA sequence encoding for wild-type (i.e., endogenous) human c-Raf is depicted in the nucleic acid sequence shown in FIG. 7 (SEQ ID NO.: 1, nucleotides 130...2076) with the encoded translated amino acid residue sequence for the Raf protein depicted in FIG. 8 (SEQ ID NO.: 2).

The present invention describes two categories of c-Raf function to modulate angiogenesis. As previously discussed, one category contains Raf molecules that increase angiogenesis and, thus, are considered to be active proteins. Wild-type Raf along with various mutations are shown in the present invention to induce angiogenesis.

One preferred mutation of wild type c-Raf which functions in this context with respect to its ability to induce blood vessel growth and therefore increase tumor weight *in vivo* is the Raf mutant construct in which only the amino acid residues 306-648 of

Raf (Raf 306-648) are expressed. This construct lacks the entire regulatory kinase domain and is therefore referred to as a constitutively active Raf protein.

5 Mutations in Raf have also been shown to have the opposite modulatory effect on angiogenesis, inhibiting angiogenesis instead of stimulating it. Such mutations are referred to as inactive Raf mutations. Proteins having mutations that confer this inhibitory activity are also referred to as dominant negative Raf proteins in that they inhibit neovascularization, including that resulting from endogenous activity of Raf as well as enhanced Raf activity resulting from growth factor stimulation. Thus, certain mutations of wild type c-Raf of the present invention can also function as a dominant
10 negative with respect to their ability to block blood vessel growth, and for example, therefore decrease tumor weight *in vivo*.

An exemplary inhibitory Raf construct is the Raf mutation in which the lysine amino acid residue 375 is mutated into any other amino acid, preferably a methionine (i.e., Raf K375M). This point mutation in the kinase domain prevents ATP binding
15 and also blocks kinase-dependent Raf functions related to vascular cell and tumor cell signaling and proliferation. Another inhibitory Raf mutant would comprise amino acid residues 1-305 in the form of a truncated Raf protein (i.e., Raf 1-305), which lacks the kinase domain.

With respect to the point mutations, any mutation resulting in the desired
20 inhibitory or stimulatory activity is contemplated for use in this invention. Fusion protein constructs combining the desired Raf protein (mutation or fragment thereof) with expressed amino acid tags, antigenic epitopes, fluorescent protein, or other such protein or peptides are also contemplated, so long as the desired modulating effect of the Raf protein is intact.

25 To produce the desired c-Raf mutations in the cDNA, standard site-directed mutagenesis procedures familiar to one of ordinary skill in the art were utilized. PCR primers designed to incorporate the desired mutations were also designed with restriction sites to facilitate subsequent cloning steps. Entire segments of Raf encoding nucleic acid sequences are deleted from the nucleic acid constructs through

PCR amplification techniques based on the known cDNA sequences of chicken, human and the like homologs of Raf and subsequent formation of new constructs.

Specifically, the wild-type Raf cDNA sequence shown in FIG. 7 was modified in several ways to construct Raf mutants to demonstrate the principles of the present invention. These mutants were inserted into the retrovirus expression system described herein.

A first mutant Raf, designated Raf K375M, was constructed in wild-type human Raf in which lysine at amino acid residue position 375 was substituted by a methionine. Raf K375M is an "inactive" Raf protein as defined herein.

A second mutant Raf, designated Raf 306-648, was constructed in wild-type human Raf in which the amino terminal portion was deleted, leaving the truncated carboxy terminal residues 306-648. Raf 306-648 is an "active" Raf protein as defined herein.

A third mutant Raf, designated Raf 1-305, is constructed in wild type human Raf in which the carboxy terminal portion was deleted, leaving the truncated amino terminal residues 1-305. Raf 1-305 is an "inactive" Raf protein as defined herein.

Alternative expression vectors for use in the expressing the Raf or Ras proteins of the present invention also include adenoviral vectors as described in US Patent No. 4,797,368, No. 5,173,414, No. 5,436,146, No. 5,589,377, and No. 5,670,488.

Alternative methods for the delivery of the Raf or Ras modulatory proteins include delivery of the Raf or Ras cDNA with a non-viral vector system as described in US Patent No. 5,675,954 and delivery of the cDNA itself as naked DNA as described in US Patent No. 5,589,466. Delivery of constructs of this invention is also not limited to topical application of a viral vector, viral vector preparations are also injected intravenously for systemic delivery into the vascular bed, or can be injected subcutaneously, intratissue, and the like. These vectors are also targetable to sites of increased neovascularization by localized injection of a tumor, as an example.

In vitro expressed proteins are also contemplated for delivery thereof following expression and purification of the selected Raf or Ras protein by methods useful for

delivery of proteins or polypeptides. One such method includes liposome delivery systems, such as described in US Patent No. 4,356,167, No. 5,580,575, No. 5,542,935 and No. 5,643,599. Other vector and protein delivery systems are well known to those of ordinary skill in the art for use in the expression and/or delivery of the Raf or Ras proteins of the present invention.

2. Human Tumor Model

To demonstrate the efficacy of the present invention, human tumor cells were implanted subcutaneously onto the flank of athymic mice, and allowed to grow to about 100 mm³. In this xenograft model, the murine endothelial cells in the tissue surrounding the implant form vasculature that grow into the growing human tumor in response to the normal angiogenic signals, and the tumor becomes vascularized. Thus, the microvessels are formed by murine endothelial cells, whereas the tumor tissue itself comprises human cells.

3. Retrovirus Delivery Vector Infects Mouse Lineage Cells, Not Human Tumor Cells

The retrovirus expression vector system of Clonetech was used to construct ecotrophic retrovirus which contain the constructs of Raf described herein. To demonstrate the tissue specificity of the infecting retrovirus, a retrovirus expression vector construct which expresses b-galactosidase was packaged using ecotrophic packaging cells as described in the legend to FIG. 1.

Mouse 3T3, mouse endothelial cells, human epithelial adenocarcinoma LS174 cells and human melanoma M21 cells were cultured *in vitro*, and were each exposed to the ecotrophically packaged retrovirus. Only the murine cells express detectable b-galactosidase, indicating that only murine cells are infected by ecotrophically packaged retrovirus in this expression system.

4. Inactive Raf Kinase Disrupts Raf Kinase Activity In Vitro

To demonstrate the cellular effects of inactive Raf kinase, an *in vitro* model using mouse endothelial cells induced by bFGF was used. The normal induction of Raf activity by bFGF administration to mouse endothelial cells was blocked when those cells were first infected by a retroviral construct which expressed the inactive Raf K375M kinase construct as described in the legend to FIG. 2. The data in FIG. 2 shows that the amount of Raf kinase activity is substantially reduced when cells are first infected by the vector which expresses an inactive Raf kinase.

5. Inactive Raf Kinase Disrupts Angiogenesis In Vivo

Using an *in vivo* murine subcutaneous model for angiogenesis, the effects of inactive Raf kinase were studied. To that end, angiogenesis was induced in a mouse by injection of bFGF either with or without cells expressing retrovirus that produces the inactive Raf K375M kinase protein as described in the legend to FIG. 3. As shown in FIG. 3, the presence of inactive Raf kinase substantially reduced the angiogenic index.

6. Active Raf Kinase Induces Angiogenesis In Vivo

Using the murine subcutaneous model for angiogenesis, the effects of active Raf kinase were studied. To that end, angiogenesis was induced by injection of cells expressing retrovirus that produced the active Raf 306-648 kinase as described in the legend to FIG. 4. As shown in FIG. 4, mutationally active Raf kinase induces angiogenesis *in vivo*.

7. Inactive Raf Kinase Induces Apoptosis

Using the mouse xenograft model described above, the *in vivo* effects of inactive Raf were studied. To that end, the model was established as described in the legend to FIG. 5 by injection of 1.5 million human adenocarcinoma LS174 cells. Following establishment of a tumor mass of about 100 mm³, retrovirus expressing the inactive Raf K375M kinase were injected into the tumor mass, and

immunohistochemistry was performed on sections of the tumor mass after 48 hours. The results shown in FIG. 5 (Flag tag) indicate that the retrovirus infection was endothelial specific, and further shows via the vWF stain that the endothelial cells colocalized to the retrovirus infection. The merge of the staining data shows that the endothelial cells, the virus infection and the occurrence of apoptosis all colocalized, indicating that the virus delivery of the inactive Raf protein is endothelial specific and that inactive Raf induces apoptosis.

8. Inactive Raf Kinase Induces Tumor Regression

Using the mouse xenograft model described above, the *in vivo* effects of inactive Raf on tumor regression were studied. To that end, the model was established as described in the legend to FIG. 6, and the inactive Raf K375M kinase was provided as virus supernate or virus-expressing cells as indicated. The established tumor was seen to rapidly regress upon introduction of inactive Raf kinase.

9. Angiogenesis is Dependent on Activation of the Ras-Raf-MEK-ERK Pathway

To determine the interaction of growth factor receptor and integrin receptor ligation and activation on the activation of the mitogen-activated protein kinase (MAPK)/extracellular signal-regulated kinase (ERK) cascade that is involved in modulating angiogenesis, the following studies in Examples 9-11 were performed. Activation of the MAPK cascade by integrin-mediated cell adhesion has been investigated by a number of laboratories as reviewed by Aplin et al., Pharmacol. Rev., 50:197-263 (1998). The hierarchical ERK cascade originates at the cell membrane with receptors for mitogens and growth factors which recruits the small guanosine triphosphate (GTPase) Ras which then activates Raf, a protein kinase, by binding to Raf and recruiting it to the membrane, where it is activated in a yet undetermined mechanism. Activated Raf then phosphorylates and activates MEK (MAPK/ERK kinase). MEK, then, phosphorylates and activates ERK1 and ERK2 which then translocate to the nucleus and transactivate transcription factors to effect growth,

differentiation or mitosis through altered gene expression. (See, Tibbles et al., Cell Mol. Life Sci., 55:1230-1254 (1999)).

5 The upstream regulation of Ras in activation of Raf that is mediated by growth factor and/or integrin signaling is the subject of current studies but the mechanisms of signaling are still not completely understood. (See, Stewart et al., J. Biol. Chem., 275:8854-8862 (2000); Howe et al., J. Biol. Chem., 273:27268-27274 (1998)). However, and more importantly, the activation of the Ras-Raf-MEK-ERK cascade through cell membrane receptor signaling resulting in modulation of angiogenesis has not been described before the present invention.

10 A. Ras is Induced by Exposure to bFGF

Therefore, to first assess whether angiogenesis was dependent on the Ras-Raf-MEK-ERK pathway, Ras activity was measured in chick chorioallantoic membrane (CAM) lysates exposed to bFGF as determined by a Ras pulldown assay.

15 Angiogenesis can be induced on the CAM after normal embryonic angiogenesis has resulted in the formation of mature blood vessels. Angiogenesis has been shown to be induced in response to specific cytokines or tumor fragments as described by Leibovich et al., Nature, 329:630 (1987) and Ausprunk et al., Am. J. Pathol., 79:597 (1975). CAMs were prepared from chick embryos for subsequent induction of angiogenesis and inhibition thereof. Ten day old chick embryos were
20 obtained from McIntyre Poultry (Lakeside, CA) and incubated at 37°C with 60% humidity. A small hole was made through the shell at the end of the egg directly over the air sac with the use of a small crafts drill (Dremel, Division of Emerson Electric Co. Racine WI). A second hole was drilled on the broad side of the egg in a region devoid of embryonic blood vessels determined previously by candling the egg.
25 Negative pressure was applied to the original hole, which resulted in the CAM (chorioallantoic membrane) pulling away from the shell membrane and creating a false air sac over the CAM. A 1.0 centimeter (cm) x 1.0 cm square window was cut

through the shell over the dropped CAM with the use of a small model grinding wheel (Dremel). The small window allowed direct access to the underlying CAM.

The resultant CAM preparation was then used at 10 days of embryogenesis where angiogenesis has subsided. The latter preparation was, thus, used in this invention for inducing renewed angiogenesis in response to cytokine treatment or tumor contact, where necessary, as described below.

1) Angiogenesis Induced by Growth Factors

Angiogenesis has been shown to be induced by cytokines or growth factors. Angiogenesis was induced by placing a 5 millimeter (mm) X 5 mm Whatman filter disk (Whatman Filter paper No.1) saturated with Hanks Balanced Salt Solution (HBSS, GIBCO, Grand Island, NY) or HBSS containing recombinant basic fibroblast growth factor (bFGF) or vascular endothelial cell growth factor (VEGF) (Genzyme, Cambridge, MA) on the CAM of either a 9 or 10 day chick embryo in a region devoid of blood vessels and the windows were latter sealed with tape. Other growth factors are also effective at inducing blood vessel growth. For assays where inhibition of angiogenesis is evaluated with intravenous injections of antagonists, such as LM609 monoclonal antibody, angiogenesis is first induced with bFGF or VEGF in fibroblast growth medium, and then inhibitors are administered as described in Example 10. Angiogenesis was monitored by photomicroscopy after 72 hours.

CAMs from 10-day old chick embryos were stimulated topically with filter disks saturated with either PBS or 30 nanograms (ng) of bFGF. After 5 minutes, CAM tissue was resected, homogenized in lysis buffer, and Ras activity was then determined by its capacity to be precipitated by a GST fusion peptide encoding the Ras binding domain of Raf. Because only active Ras binds Raf, a recombinant protein was generated consisting of the Ras binding domain of Raf conjugated to glutathione-S-transferase (GST). In turn GST was conjugated to sepharose beads enabling the precipitation of active Ras from a tissue lysate.

The results are shown in FIG. 9 where Ras activity was elevated in CAM lysates exposed to bFGF as determined by a Ras pulldown assay. Thus, Ras is induced with exposure to bFGF in the CAM. The role of Ras in the formation of angiogenic blood vessels in the CAM is further assessed as described in Example 10.

5 B. Ras is Necessary for Angiogenesis

To then determine whether angiogenesis was dependent on the activation of Ras in the CAM preparation, the CAM was exposed to RCAS retroviral preparations for expression of a dominant negative Ras mutant, S17N Ras, in combination with bFGF activation of Ras as described below. This mutant has been
10 shown to bind GDP with preferential affinity over GTP, thereby providing the mutant to inhibit endogenous Ras activation by sequestering Ras-GEFs. Thus, use of the mutant in the CAM angiogenesis model provides a method to assess the role of Ras in angiogenesis.

The S17N Ras mutant is created from the wild -type human Ras (wt H-Ras)
15 sequence by standard site directed mutagenesis procedures as previously described substituting the encoding triplet for a serine (S) residue at position 17 with a codon for encoding an asparagine (N). Such mutants have been described by others, for example, by Stewart et al., J. Biol. Chem., 275:8854-8862 (2000).

To prepare the retroviral construct of the dominant negative expression
20 construct, such mutagenesis was performed on the wt H-Ras, where the nucleic acid sequence encoding it is shown in FIG. 10 (SEQ ID NO.: 3). FIG. 11 (SEQ ID NO.: 4) depicts the amino acid residue sequence encoded by the cDNA nucleotide sequence of wild-type human Ras (wt H-Ras) shown in FIG. 10. To produce the desired mutations in the wt H-Ras cDNA to make S17N Ras as well as those described below, standard
25 site-directed mutagenesis procedures familiar to one of ordinary skill in the art were utilized. PCR primers designed to incorporate the desired mutations were also designed with restriction sites to facilitate subsequent cloning steps. Entire segments of Ras encoding nucleic acid sequences can be deleted from the nucleic acid constructs

through PCR amplification techniques based on the known cDNA sequences of chicken, human and the like homologs of Ras and subsequent formation of new constructs. All mutant constructs constructed by PCR were also sequenced by PCR to confirm predicted DNA sequence of clones.

5 The resultant mutated Ras sequence was then prepared as an retroviral expression vector construct as described herein. One preferred expression construct for use in the present invention is the RCAS(A) construct. This expression vector is based on a series of replication competent avian sarcoma viruses with an enhanced Bryan polymerase (BP) for improved titre, and is specific for the A type envelope glycoprotein expressed on normal avian cells (Reviewed in *Methods in Cell Biology*, 10 52:179-214 (1997); see also, Hughes et al., *J. Virol.* 61:3004-3012 (1987); Fekete & Cepko, *Mol. Cellular Biol.* 13:2604-2613 (1993); Itoh et al., *Development* 122:291-300 (1996); and Stott et al., *BioTechniques* 24:660-666 (1998)). The complete sequence of RCAS(A), referred to herein as RCAS, is known to one of ordinary skill 15 in the art and available on databases.

Five micrograms (ug) of RCAS constructs prepared were then transfected into the chicken immortalized fibroblast line, DF-1 (gift of Doug Foster, U. of Minn.). This cell line as well as primary chick embryo fibroblasts were capable of producing virus, however the DF-1 cell line produced higher titres. Viral supernatants were 20 collected from subconfluent DF-1 producer cell lines in serum free CLM media [composition: F-10 media base supplemented with DMSO, folic acid, glutamic acid, and MEM vitamin solution]. Thirty-five ml of viral supernatant were concentrated by ultracentrifugation at 4°C for 2 hours at 22,000 rpm. These concentrated viral pellets were resuspended in 1/100 the original volume in serum-free CLM media, aliquoted 25 and stored at -80°C. The titre was assessed by serial dilution of a control viral vector having a nucleotide sequence encoding green fluorescent protein (GFP), referred to as RCAS-GFP, infection on primary chick embryo fibroblasts that were incubated for 48-72 hours. The titres of viral stock that were obtained following concentration routinely exceeded 10⁸ I.u./ml.

For the CAM assay using the viral stocks, cortisone acetate soaked Whatman filter disks 6 mm in diameter were prepared in 3 mg/ml cortisone acetate for 30 minutes in 95% ethanol. The disks were dried in a laminar flow hood and then soaked on 20 µl of viral stock per disk for 10 minutes. These disks were applied to the CAM of a 10 day chick embryos and sealed with cellophane tape and incubated at 37°C for 18-24 hr. Then either mock PBS or growth factors were added at a concentration of 5 µg/ml to the CAM in a 15 microliters (ul) volume of the appropriate virus stock as an additional boost of virus to the CAM tissue. After 72 hours, the CAMs were harvested and examined for changes in the angiogenic index as determined by double blind counting of the number of branch points in the CAM underlying the disk. For kinase assays, the tissue underlying the disk was harvested in RIPA, homogenized with a motorized grinder and Raf determined as previously described in Example 4. For immunofluorescence studies, CAM tissue underlying the disks were frozen in OCT, a cryopreservative, sectioned at 4 um, fixed in acetone for 1 minute, incubated in 3% normal goat serum for 1 hour, followed by an incubation in primary rabbit antibody as described previously (Eliceiri et al., J. Cell Biol., 140:1255-1263 (1998), washed in PBS and detected with a fluorescent secondary antibody.

The results, shown in FIG. 12, graphically reveal that infection with mutant null Ras, S17N, blocked growth factor-induced angiogenesis in the CAM, but had no effect on CAMs that were not exposed to bFGF to induce angiogenesis. Therefore, Ras is necessary for bFGF-induced angiogenesis.

C. Ras Signaling Through the Raf-MEK-ERK Pathway is a Crucial Regulator of Angiogenesis

To further assess the role of Ras in the Raf-MEK-ERK pathway in modulating angiogenesis, additional H-Ras mutant proteins were used in the CAM preparation as described above, the results of which are shown below and in FIG. 13. In this context, the present invention describes two categories of Ras function that can modulate angiogenesis. As previously discussed for Raf proteins, one category

contains Ras molecules that increase angiogenesis and, thus, are considered to be active proteins. Wild-type Ras along with various mutations are shown in the present invention to induce angiogenesis.

5 One preferred mutation of wild type H-Ras which functions in this context with respect to its ability to induce blood vessel growth and therefore increase tumor weight *in vivo* is the Ras G12V, also referred to as V12, mutant having a point mutation at amino acid (aa) residue position 12 changing glycine (G) to valine (V). This mutant Ras is constitutively active.

10 Another H-Ras mutant protein that is described for the present invention as a constitutive angiogenesis activator is Ras V12S35, where the glycine at position 12 was changed to valine (V) and the threonine (T) at position 35 was changed to a serine (S), both mutations resulting in Ras V12S35. This mutated H-Ras protein has been shown to only selectively activate the Raf-MEK-ERK pathway as shown in FIG. 13A.

15 A H-Ras negative regulator of angiogenesis is Ras V12C40 mutant, where the glycine at position 12 was changed to valine (V) as in Ras V12S35 but the other mutation was at position 40 where a tyrosine residue (Y) was changed to a cysteine (C), both mutations, thus, resulting in Ras V12C40. This mutant H-Ras is known to selectively activate the P1-3 kinase (P13K as shown in FIG. 13A) pathway that activates Akt and Rac. Thus, Ras V12C40 does not function in the Raf-MEK-ERK
20 pathway and does not stimulate angiogenesis but rather would inhibit it. Proteins having mutation that confer inhibitory activity on angiogenesis are also referred to as dominant negative Ras proteins in that they inhibit neovascularization, including that resulting from endogenous activity of Ras as well as enhanced Ras activity resulting from growth factor stimulation. Thus, certain mutations of wild type H-Ras of the
25 present invention can also function as a dominant negative with respect to their ability to block blood vessel growth, and for example, therefore decrease tumor weight *in vivo*. The three H-Ras constructs and mutant proteins have been previously described by Joneson et al., Science, 271:810-812 (1996).

With respect to the point mutations, any mutation resulting in the desired inhibitory or stimulatory activity is contemplated for use in this invention. Fusion protein constructs combining the desired Ras (or Raf proteins as shown in the Examples below) (mutation or fragment thereof) with expressed amino acid tags, antigenic epitopes, fluorescent protein, or other such protein or peptides are also contemplated, so long as the desired modulating effect of the Ras protein is intact.

To evaluate the roles of the additional Ras mutant proteins in signaling pathway activation of angiogenesis, the respective retroviral expression constructs were prepared as described above. Fifteen ul of high titer RCAS (A) virus encoding the Raf-MEK-ERK activating Ras construct, Ras V12S35, or the PI3 kinase activating Ras construct, Ras V12C40, were topically applied to filter disks in a 10-day old CAM preparation and results assessed as described above for the effect of the mutant Ras proteins on angiogenesis with respect to the selective activation of signaling pathways.

FIGs. 13A and 13B illustrate schematically and graphically respectively that infection with a mutant Ras construct, Ras V12S35, which selectively activates the Ras-Raf-MEK-ERK pathway, induced angiogenesis, whereas a mutant construct, Ras V12C40, which selectively activates the PI3K pathways did not. Thus, these results confirm that Ras V12S35 protein is a angiogenesis stimulator and that Ras-mediated activation of angiogenesis occurs through activation of the Raf-MEK-ERK pathway and not via the P13K pathway utilized by the H-Ras mutant V12C40.

D. The MEK Component of the MEK-ERK Pathway is Required for Either Ras or Ras-Independent Raf Induced Angiogenesis

To further assess the separate roles of Ras and Raf in the Raf-MEK-ERK pathway in modulating angiogenesis, a Raf mutant protein, referred to as Raf-Caax, that is targeted to the plasma membrane that is known to be constitutively and enzymatically active in the absence of Ras binding was used in the CAM preparations as described herein in conjunction with a known inhibitor of MEK activation, PD98059. FIG. 14 depicts the nucleotide sequence encoding the fusion protein

Raf-caax, where the nucleotide sequence encoding the carboxy terminus of human Raf (wt H-Raf) is fused with a nucleotide sequence of encoding a 20 amino acid residue sequence of the K-ras membrane localization domain (SEQ ID NO.: 6). FIG. 15 (SEQ ID NO.: 7) depicts the amino acid residue sequence of Raf-caax, the fusion protein generated from the fusion nucleotide sequence shown in FIG. 14. The fusion protein has been described by Leever et al., Nature, 369:411-414 (1994) and Stokoe et al., Science, 264:1463-1467 (1994).

For assessing the Ras-independent Raf-induced angiogenesis along with angiogenesis induced by Raf, the MEK inhibitor, PD98059, was used in CAM preparations as described above. Virus encoding the activating Ras construct, Ras V12 (Ras G12V), prepared as described in Example 9C and the activating Raf construct, Raf-caax, were topically applied to filter disks as described in Example 9B. After 24 hours, one (1) nanomole of the MEK inhibitor, PD98059, was added to the disk. The CAMs were then evaluated as described in Example 9B and in FIG. 12. Data plotted is the mean \pm SE of 20 embryos.

FIGs. 16A-16E and FIG. 16F, respectively, pictorially and graphically illustrate that the MEK inhibitor, PD98059, blocked angiogenesis (FIGs. 16C and 16E) induced by either mutant active Ras (FIG. 16B) or Raf (FIG. 16D). Thus, both Ras and Raf induce angiogenesis through the MEK-ERK pathway. The plotted data graphically depicts the results of the photographs of the individual treated CAMs.

10. Angiogenesis induced by Raf, but not Ras, is Refractory to Inhibition by Integrin Blockade

To determine how integrin signaling activates the Ras-Raf-MEK-ERK pathway resulting in angiogenesis, CAM assays with mutant active Ras and Raf constructs were performed in the presence of $\alpha_v\beta_3$ integrin-blocking antibodies. CAMs from 10-day old chick embryos were stimulated as described in FIGs. 9 and 12 with filter disks saturated with either PBS (control), bFGF, the RCAS(A) retroviral constructs G12V-Ras or Raf-caax. LM609, a monoclonal antibody to integrin $\alpha_v\beta_3$,

was intravenously delivered after 24 hours and angiogenesis was assessed by vessel branch point analysis after 72 hours. Representative CAMs are shown in the inset. Data is the mean \pm SE of 20 embryos.

FIGs. 17A-17F and FIG. 17G, respectively, pictorially and graphically illustrate that angiogenesis induced by Raf, but not Ras, was refractory to inhibition by integrin blockade. Infection with both mutant active Ras and Raf constructs induced pronounced angiogenesis as shown respectively in FIGs. 17B and 17C, but only Ras-induced angiogenesis was inhibited by $\alpha_v\beta_3$ integrin-blocking antibodies as shown in FIG. 17E. Since the Raf construct used in the assay is Ras-independent, the lack of integrin inhibition of Raf-induced angiogenesis indicates that integrin signaling occurs at or before Ras-mediated activation of Raf. The plotted data graphically depicts the results of the photographs of the individual treated CAMs.

11. Regulation of the Ras-Raf-MEK-ERK Pathway by Focal Adhesion Kinase

To determine the role of growth factor receptor activation of the Ras-Raf-MEK-ERK angiogenesis pathway, CAM angiogenesis assays were performed as described above with either Ras V12 or Raf-caax expressed proteins in the presence of a mutant null focal adhesion kinase, referred to as FRNK, which is an inactive focal adhesion kinase.

RCAS(A) viruses encoding Ras V12 or Raf-caax, prepared as described above, were topically applied as described in Example 9B (FIG. 12) along with RCAS(B) virus encoding FAK-related-null-kinase (FRNK) to the CAM filter disk. Data is the mean \pm SE of 20 embryos.

The results are shown in FIGs. 18A-18D and 18E. FIGs. 18A-18D pictorially illustrate that co-infection of CAMs with a mutant null focal adhesion kinase, FRNK, blocked Ras, but not Raf-induced angiogenesis, as indicated by a paucity of blood vessels in FIG. 18B as compared to untreated Ras (FIG. 18A), untreated Raf (FIG. 18C) and FRNK-treated Raf (FIG. 18 D). The plotted data graphically depicts the results of the photographs of the individual treated CAMs.

The data in the CAM assay was confirmed in the murine subcutaneous angiogenesis model, prepared as previously described. Angiogenesis was induced by injecting 250 ul of ice-cold, growth factor-reduced matrigel containing either 400 ng/ml bFGF or Moloney retrovirus expressing packaging cells expressing the described gene, subcutaneously in the mouse flank. FRNK retrovirus was added to matrigel as high titer virus packaged with the vsv.g coat protein. Five days later, endothelial-specific FITC-conjugated Bandeiraea Simplicifolia B5 lectin was injected via the tail vein and allowed to circulate. Angiogenesis was then quantitated by removing, extracting, and assaying the angiogenic tissue for fluorescent content.

FIGs. 19A and 19B-19G, respectively, graphically and pictorially, illustrate that FRNK blocked bFGF and Ras-, but not Raf, -induced angiogenesis in a murine subcutaneous angiogenesis model.

To verify the level at which kinase activation occurs in the Ras-Raf-MEK-ERK pathway, CAMs were co-infected with a retrovirus expressing FRNK, the mutant null focal adhesion kinase, with either Ras G12V or Raf-caax. CAMs were treated as described in FIG. 18 with the exception that after 24 hours the angiogenic tissue was resected, solubilized, Raf immunoprecipitated, and Raf activity assessed by its capacity to phosphorylate kinase-dead MEK. FIGs. 20A and 20B illustrate that co-infection of CAMs with a mutant null focal adhesion kinase, FRNK, blocked Ras-induced activation of Raf. FIG. 20A shows the immunoprecipitated active versus total Raf proteins assayed under each of the combinations above the results. FIG. 20B graphically plots the results of the active Raf determinations under those conditions. Thus, FRNK does not directly inhibit the activity of Raf but rather inhibits the activation of Raf by Ras.

12. Discussion

The above studies indicates that Raf kinase is necessary and sufficient for angiogenesis *in vivo*. Further, targeting of mutationally inactive Raf kinase to growing blood vessels induces local endothelial apoptosis. The same targeting also suppresses

angiogenesis which results in suppression and even regression of pre-existing human tumors.

The retroviral delivery of a gene encoding mutationally inactive forms of Raf kinase (Raf K375M) demonstrated a substantial impact on tumor angiogenesis *in vivo*.
5 Importantly, the retroviral vector used specifically infects proliferating cells of murine lineage. Therefore, only the vascular compartment of human tumor xenografts was infected (FIGs. 1 and 4). Delivery of inactive Raf K375M kinase was found to suppress growth factor-induced Raf kinase activity *in vitro* and block growth factor-induced angiogenesis *in vivo* (FIGs. 2 & 3). In contrast, retroviral delivery of a
10 mutationally active form of Raf kinase (Raf 306-648) was sufficient to induce angiogenesis *in vivo* (FIG. 4). Furthermore, the delivery of virus expressing inactive Raf kinase to the tumor in mice was found to induce apoptosis in a endothelial-specific manner (FIG. 5). Finally, animals inoculated with human tumors and then treated with the virus expressing inactive Raf experienced a rapid tumor regression which was
15 maintained throughout the time-course of the experiment (FIG. 6). Therefore, Raf kinase is both sufficient and necessary for angiogenesis and targeting this kinase can suppress angiogenesis and obviate angiogenesis-dependent disease.

As a result of the foregoing angiogenesis assays in mouse and chicken as described in Examples 9-11, depicted in FIGs. 9, 12, 13, and 16-20, the present
20 invention provides angiogenesis activator proteins in Raf-caax, Ras G12V Ras, and Ras V12S35 and angiogenesis inhibitor proteins in Ras S17N and Ras V12C40. Furthermore, the studies provide the basis for understanding the Ras-mediated activation of Raf in the Ras-Raf-MEK-ERK pathway identifying that Ras is necessary for activation of Raf but integrin-mediated signaling interacts at of before Raf
25 activation but not downstream thereof.

While the foregoing written specification is sufficient to enable one skilled in the art to practice the invention, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

What Is Claimed Is:

1. An article of manufacture comprising packaging material and a pharmaceutical composition contained within said packaging material, wherein said pharmaceutical composition is capable of modulating angiogenesis in a tissue associated with a disease condition, wherein said packaging material comprises a label which indicates that said pharmaceutical composition can be used for treating disease conditions by modulating angiogenesis, and wherein said pharmaceutical composition comprises a Raf protein or an oligonucleotide having a nucleotide sequence capable of expressing said protein.
2. The article of manufacture of claim 1 wherein said Raf protein is an active Raf protein and said modulating potentiates angiogenesis.
3. The article of manufacture of claim 2 wherein said active Raf protein is wild-type Raf.
4. The article of manufacture of claim 3 wherein said active Raf protein is a fusion protein.
5. The article of manufacture of claim 4 wherein said active Raf fusion protein is Raf-caax.
6. The article of manufacture of claim 2 wherein said tissue has poor circulation.
7. The article of manufacture of claim 1 wherein said Raf protein is an inactive Raf protein and said modulating inhibits angiogenesis.
8. The article of manufacture of claim 7 wherein said inactive Raf protein has a mutation at residue 375 such that the amino acid at position 375 is not lysine.
9. The article of manufacture of claim 7 wherein said tissue is inflamed and said condition is arthritis or rheumatoid arthritis.
10. The article of manufacture of claim 7 wherein said tissue is a solid tumor or solid tumor metastasis.
11. The article of manufacture of claim 10 wherein said administering is conducted in conjunction with chemotherapy.

12. The article of manufacture of claim 7 wherein said tissue is retinal tissue and said condition is retinopathy, diabetic retinopathy or macular degeneration.

13. The article of manufacture of claim 7 wherein said tissue is at the site of coronary angioplasty and said condition is restenosis.

5 14. The article of manufacture of claim 1 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

15. The article of manufacture of claim 1 wherein said administering comprises a single dose intravenously.

10 16. The article of manufacture of claim 1 wherein said pharmaceutical composition further comprises a liposome.

17. The article of manufacture of claim 1 wherein said pharmaceutical composition comprises a viral expression vector capable of expressing said nucleotide sequence.

15 18. The article of manufacture of claim 1 wherein said pharmaceutical composition comprises an non-viral expression vector capable of expressing said nucleotide sequence.

20 19. A method for modulating angiogenesis in a tissue associated with a disease condition comprising administering to said tissue an angiogenesis modulating amount of a pharmaceutical composition comprising a Raf protein or a nucleotide sequence capable of expressing said protein.

20. The method of claim 19 wherein said Raf protein is an active Raf protein and said modulating potentiates angiogenesis.

25 21. The method of claim 20 wherein said active Raf protein is wild-type Raf.

22. The method of claim 21 wherein said active Raf protein is a fusion protein.

23. The method of claim 22 wherein said active Raf fusion protein is Raf-caax.

24. The method of claim 20 wherein said tissue has abnormal circulation.

25. The method of claim 19 wherein said Raf protein is an inactive Raf protein and said modulating inhibits angiogenesis.

5 26. The method of claim 25 wherein said inactive Raf protein has a mutation at residue 375 such that the amino acid at position 375 is not lysine.

27. The method of claim 25 wherein said tissue is inflamed and said condition is arthritis or rheumatoid arthritis.

28. The method of claim 25 wherein said tissue is a solid tumor or solid tumor metastasis.

10 29. The method of claim 28 wherein said administering is conducted in conjunction with chemotherapy.

30. The method of claim 25 wherein said tissue is retinal tissue and said condition is retinopathy, diabetic retinopathy or macular degeneration.

15 31. The method of claim 25 wherein said tissue is at the site of coronary angioplasty and said tissue is at risk for restenosis.

32. The method of claim 19 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

33. The method of claim 19 wherein said administering comprises a single dose intravenously.

20 34. The method of claim 19 wherein said pharmaceutical composition further comprises a liposome.

35. The method of claim 19 wherein said pharmaceutical composition comprises an retroviral expression vector capable of expressing said nucleotide sequence.

25 36. The method of claim 19 wherein said pharmaceutical composition comprises an non-viral expression vector capable of expressing said nucleotide sequence.

37. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid,

said nucleic acid having a nucleic acid segment encoding for a Raf protein, said Raf protein having kinase activity, and a pharmaceutically acceptable carrier or excipient.

5 38. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic acid, said nucleic acid having a nucleic acid segment encoding for a Raf protein, said Raf protein having kinase activity, and a pharmaceutically acceptable carrier or excipient.

10 39. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid, said nucleic acid having a nucleic acid segment encoding for a Raf protein, said Raf protein having no kinase activity, and a pharmaceutically acceptable carrier or excipient.

15 40. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic acid, said nucleic acid having a nucleic acid segment encoding for a Raf protein, said Raf protein having no kinase activity, and a pharmaceutically acceptable carrier or excipient.

20 41. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a therapeutic amount of a Raf protein, said Raf protein having kinase activity, and a pharmaceutically acceptable carrier or excipient.

42. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a therapeutic amount of a Raf protein, said Raf protein having no kinase activity, and a pharmaceutically acceptable carrier or excipient.

25 43. An article of manufacture comprising packaging material and a pharmaceutical composition contained within said packaging material, wherein said pharmaceutical composition is capable of modulating angiogenesis in a tissue associated with a disease condition, wherein said packaging material comprises a label which indicates that said pharmaceutical composition can be used for treating disease conditions by modulating angiogenesis, and wherein said pharmaceutical composition

comprises a Ras protein or an oligonucleotide having a nucleotide sequence capable of expressing said protein.

44. An article of manufacture according to claim 43 wherein said Ras protein or oligonucleotide encoding said protein is an inhibitory Ras protein.

5 45. An article of manufacture of claim 44, wherein said inhibitory Ras is Ras V12C40.

46. An article of manufacture of claim 44, wherein said inhibitory Ras is Ras S17N.

10 47. An article of manufacture according to claim 43 wherein said Ras protein or oligonucleotide encoding said protein is a stimulatory Ras protein.

48. An article of manufacture of claim 47, wherein said stimulatory Ras is Ras G12V.

49. An article of manufacture of claim 47, wherein said stimulatory Ras is Ras V12S35.

15 50. A method for modulating angiogenesis in a tissue associated with a disease condition comprising administering to said tissue an angiogenesis modulating amount of a pharmaceutical composition comprising a Ras protein or a nucleotide sequence encoding for said protein.

20 51. The method according to claim 50 wherein said Ras protein or oligonucleotide encoding for said protein is an inactive Ras protein.

52. The method of claim 51, wherein said inactive Ras is Ras V12C40.

53. The method of claim 51, wherein said inactive Ras is Ras S17N.

54. The method according to claim 50 wherein said Ras protein or oligonucleotide encoding for said protein is an active Ras protein.

25 55. The method of claim 54, wherein said active Ras is Ras GV12.

56. The method of claim 54, wherein said active Ras is Ras V12S35.

57. A pharmaceutical composition for modulating angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid, said nucleic acid having a nucleic acid segment encoding for a Ras protein, said Ras

protein having angiogenesis modulating activity, in a pharmaceutically acceptable carrier or excipient.

5 58. A pharmaceutical composition of claim 57, for inhibiting angiogenesis in a target mammalian tissue wherein said Ras protein has angiogenesis inhibitor activity.

59. A pharmaceutical composition of claim 58, wherein said Ras protein is Ras V12C40.

60. A pharmaceutical composition of claim 58, wherein said Ras protein is Ras S17N.

10 61. A pharmaceutical composition of claim 57, wherein said Ras protein has angiogenesis activating activity.

62. A pharmaceutical composition of claim 61, wherein said Ras protein is Ras G12V.

15 63. A pharmaceutical composition of claim 61, wherein said Ras protein is Ras V12S35.

20 64. A pharmaceutical composition for modulating angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic acid, said nucleic acid having a nucleic acid segment encoding for a Ras protein, and said Ras protein having angiogenesis modulating activity, and a pharmaceutically acceptable carrier or excipient.

25 65. A method for modulating angiogenesis in a tissue associated with a disease condition comprising administering to said tissue an angiogenesis modulating amount of a pharmaceutical composition comprising a Raf protein or a nucleotide sequence capable of expressing said protein, and a Ras protein or a nucleotide sequence capable of expressing said protein.

66. A method of claim 65 wherein said modulation is an inhibition of angiogenesis, and at least one of said Raf and Ras proteins are inactive.

67. A method of claim 65 wherein said modulation is an stimulation of angiogenesis, and at least one of said Raf and Ras proteins are active.

ABSTRACT

5 The present invention describes methods for modulating angiogenesis in tissues using Raf and/or Ras protein, modified Raf or Ras protein, and nucleic acids encoding for such. Particularly the invention describes methods for inhibiting angiogenesis using an inactive Raf and/or Ras protein, or nucleic acids encoding therefor, or for potentiating angiogenesis using an active Raf and/or Ras protein, or nucleic acids encoding therefor. The invention also describes the use of gene delivery systems for providing nucleic acids encoding for the Raf or Ras protein, or modified forms thereof.




FIGURE 1






FIGURE 2

A



B





FIGURE 3

A



B




FIGURE 4


[illegible]

Endothelial cells

↙

infected cells

apoptotic cells



infected, apoptotic ECs

FIGURE 5





FIGURE 6

HUMAN c-RAF cDNA

ccgaatgtgaccgctcccgtccctcaccgcccggggaggaggagcggcgagaagctccgcccgaacgacaggacg
 ttggggcggcctggctccctcaggtttaagaattgtttaagctgcatcaatggagcacatacaggagcttgaagacga
 tcagcaatgggtttgattcaaatgacggtgttgatggctccagctgcatctctctacaatagttcagcagttggc
 tatcagcggcgggcatcagatgatggcaactcacagatccttctaagacaagcaacactatccgtgttttctgccga
 caagcaagaacagtggtcaatgtgcgaatggatgagcttgcagtgactgccttatgaaagcactcaagggtgaggggccc
 tgcaaccagagtgtgtgcagtggtcagacttctccacgaacacaaaggtaaaaagcacgcttagattggaatactgat
 gctgctgtttgattggagaagaactcaagtagatttctggtatgttccctcacaacacacaacttctcggaa
 gacgttctgaagcttgccttctgtgacatctgtcagaaattctgtcaatggatttcgatgtcagacttgggtaca
 aatttcagcagctgtagcaccaaagtacactatgtgtgtgactggagtaacatcagacaactctattgtttcca
 aattccactattggtgatagtgaggtcccagcactaccttcttactatgctgctgatgcgagagtctgtttccaggt
 gcctgttagttctcagcacagatattctacacctcacgcttcccttaacacctccagtcctctcatcgaagggtccc
 tctccagaggcagaggtcagatccacacctaattgtccacatggtcagcaccacgctgcctgtgacagcaggatgatt
 gaggatgcaattcgaagtcacagcgaatcagcctcaccttcagccctgtccagtagccccaacaatctgagcccaacagg
 ctggtcacagccgaaaacccccgtgccagcacaaagagagcgggaccagtatctgggaccaggagaaaaacaaatta
 ggctcgtggacagagagattcaagctattattgggaaatagaagccagtgaagtgtgctgtccactcggattgggtca
 ggctcttttgaactgtttataagggttaaattggcacggagatgttcagtaaaagatcctaaagggttgcgacccaacccc
 agagcaattccaggccttcagggaatgaggtggtgttctcgcacaaacacggcatgtgaacattctgttttcattgggt
 acatgacaaaggacaacctggcaattgtgacctggtggtgagggcagcagcctctacaaacacctgcatgtccaggag
 accaagtttcagatgttccagctaatgacattgcccggcagacggctcagggaatggactatttgcagcaagaacat
 catccatagagacatgaaatccaacaatatatttctccatgaaggcttaacagtgaatggagattttgtttggcaa
 cagtaaaagtcacgctggagtggttctcagcaggttgaaacaacctactggctctgtcctctggatggccccagagggtatc
 cgaatgcaggataacaacccattcagttccagtcggatgtctactcctatggcatcgtattgtatgaactgatgacggg
 ggagcttcttattctcacatcaacaaccgagatcagatcatcttcattggtgggcccagggatattgcctccccagatcta
 gtaagctatataagaactgccccaaagcaatgaagaggctgtgactgtgaagaaagtaagggaagagagcct
 cttttccccagatcctgtcttccattgagctgtcctaacactctctaccgaagatcaaccggagcgttccgagccatc
 cttgcatcgggcagcccacactgaggtatatcaatgcttgcacgctgaccacgtcccagggtgcctgtcttctagtga
 ctttgacactgttctcaggctgccaggggaggaggagaagccagcaggcaccacttttctgtcctcttctccagaggca
 gaacacatgtttcagagaagctctgctaaggacctcttagactgctcacagggccttaactcatgttgccttcttct
 tatccctttgggcccctgggagaaggaagccatttgcagtgtggtgtgtcctgtcctccccacattccccatgctcaa
 ggcccagccttctgtagatgcgcaagtggatgttgatgtagtacaaaaagcaggggcccagcccagctgttggctaca
 tgagtatttagaggaagtaaggttagcaggcagtcagccctgatgtggagacacatgggattttgaaatcagcttctgg
 aggaatgcatgtcacaggcgggactttctcagagagtgtgtgcagcgcagacattttgcacataaggccacaaacagcc
 caggactgccgagactctggcccggcgaaggagcctgcttgggtactatggaactttcttaggggacacgtcctcctt
 cacagcttctaagggtgtccagtgcatgggatggtttccaggcaaggcactcggccaatccgcattcagccctcag
 gagcagcttccatcatgtgaattttgttctccaggagctgccctatggggcgggccgcaggggccagcctgtttctct
 aacaaacaaacaaacaaacagccttgtttctctagtcatcatgtgtatacaagggaagccaggaatacaggtttcttg
 atgatttgggttttaattttgtttttattgcacctgacaaaatacagttatctgatgtcctcaattatgttatttaa
 taaaataaataaattt

FIGURE 7

[illegible]

FIGURE 8




FIGURE 9

atgacggaatataagctggtggtggtggcgccggcggtgtgggcaagagtgcgctgaccatccagctgatccagaacca
 ttttgtggacgaatacgaacccactatagaggattcctaccggaagcaggtggtcattgatggggagacgtgcctgttgg
 acatcctggataccgcccggccaggaggagtacagcgccatgcccggaccagtacatgcgcaccggggagggttctgtgt
 gtgtttgccatcaacaacaccaagcttttgaggacatccaccagtacaggagcagatcaaacgggtgaaggactcggga
 tgacgtgcccattggtgctggtgggggaacaagtgtgaccggctgcacgactgtggaatctcggcagggtcaggacctcg
 ccgaagctacggcatcccctacatcgagacctcggccaagaccggcgaggagtggaggatgccttctacacgttgggtg
 cgtgagatccggcagcacaaagctgcggaagctgaaccctcctgatgagagtggccccggctgcatgagctgcaagtgtgt
 gctctcctga

FIGURE 10

MTEYKLVVVGAGGVGKSALTIQLIQNHFVDEYDPTIEDSY
 RKQVVIDGETCLLDILDITAGQEEYSAMRDQYMRTGEGFLC
 VFAINNTKSFEDIHQYREQIKRVKDSDDVPMVLVG NKCDL
 AARTVESRQAQDLARSYGIPYIETSAKTRQGVEDAFYTLV
 REIRQHKLRKLNPPDESGPGCMSCKCVLS

FIGURE 11




FIGURE 12

Ras Effector Mutants




FIGURE 13A




FIGURE 13B

atggagcacatacagggagcttggagacgatcagcaatgggtttggattcaaagatgccgtgttgatggctccagctg
catctctctacaatagttcagcagtttggctatcagcgccgggcatcagatgatggcaactcacagatccttctaaga
caagcaacactatccgtgttttcttggcgaacaagcaagaacagtggtcaatgtgcgaaatggaatgagcttgcatgac
tgccttatgaaagcactcaaggtgaggggctgcaaccagagtgtgtgagtggtcagacttctcacgaacacaaagg
taaaaaagcacgcttagattggaatactgatgctgcgtctttgattggagaagaactcaagtagatttcttgatcatg
ttccctcacacacacaactttgctcggagacgttctgaagcttgcttctgtgacatctgtcagaaattctgctc
aatggattcgtatgacagacttgggtacaaattcatgagcactgtagcaccaaagtacactactatgtgtgtggactg
gagtaacatcagacaactcttattgtttccaaattccactattgggtgatgtggagtccagcactaccttctttgacta
tgcgtcgtatgcgagagtctgtttccaggaatgctgttagttctcagcacagatattctacacctcacgccttcacctt
aacacctccagtcctctcatgaaaggttccctctcccagaggcagaggtcgacatccacacctaattgtccacatgggtcag
caccacgctgcctgtggacagcaggatgattgaggatgcaattcgaaatcacagcgaatcagcctcaccttcagccctgt
ccagtagccccaacaatctgagcccaacaggctgggtcacagccgaaaacccccgtgccagcacaagagagcgggaccca
gtatctgggacccaggagaaaaacaaaattaggcctcgtggacagagagattcaagctattattgggaaatagaagccag
tgaagtgatgctgtccactcggattgggtcaggctcttttggaaactgtttataagggtaatggcacggagatgttgag
taaagatcctaaagggtgtcgacccaaccccagagcaattccaggccttcagggaatgaggtggctgttctgcgcaaaaca
cggcatgtgaacattctgcttttcatgggtacatgacaaaggacaacctggcaattgtgaccagtggtgcgagggcag
cagcctctacaaacacctgcatgtccaggagaccaagtttcagatgttcagctaatgacattgccggcagacggctc
agggaaatggactatttgcattgcaaaagaacatcatccatagagacatgaaatccaacaatatatttccatgaaggctta
acagtgaattggagattttgggttggcaacagtaaaagtcacgctggagtgttctcagcaggttgaacaacctactgg
ctctgtcctctggatggccccagaggtgatccgaatgcaggataacaacccattcagttccagtcggatgtctactcct
atggcatcgtattgtatgaactgatgacgggggagcttcttattctcacatcaacaaccgagatcagatcatcttcag
gtgggcccaggatagcctccccagatcttagtaagctatataagaactgccccaaagcaatgaagaggctggtagctga
ctgtgtgaagaaagtaaggaagagaggcctcttttcccagatcctgtcttccattgagctgtccaacactctctac
cgaagatcaaccggagcgttccgagccatccttgcatcgggcagcccacactgaggatatcaatgcttgacgctgacc
acgtccccgaggctgcctgtcttc tactcgttctcgttcttcttcttcttcttctgttctgttccacgcctagtacattc

FIGURE 14

DOT33-2025

MEHIQGAWKTISNGFGFKDAVFDGSSCISPTIVQQFGYQR
RASDDGKLTDPSTNTIRVFLPNKQRTVVNVVRNGMSLHD
CLMKALKVRGLQPECCAVFRLLEHKGKKARLDWNTDAAS
LIGEELQVDFLDHVPLTTHNFARKTFLKLAFCDICQKFL
NGFRCQTCGYKFHEHCSTKVPTMCVDWSNIRQLLLFPNST
IGDSGVPALPSLTMRRMRESVSRMPVSSQHRYSTPHAFTF
NTSSPSSEGSLSQRQRSTSTPNVHMOVSTTLPVDSRMIEDA
IRSHSESASPSALSSSPNNLSPTGWSQPKTPVPAQRERAP
VSGTQEKNKIRPRGQRDSSYYWEIEASEVMLSTRIGSGSF
GTVYKGKWHGDVAVKILKVVDPTPEQFQAFRNEVAVLRKT
RHVNILLFMGYMTKDNLAIVTQWCEGSSLYKHLHVQETKF
QMFQLIDIARQTAQGMDYLHAKNIIHRDMKSNNIFLHEGL
TVKIGDFGLATVKSRWSGSQQVEQPTGSVLWMAPEVIRMQ
DNNPFSFQSDVYSYGIVLYELMTGELPYSHINNRDQIIFM
VGRGYASPDLSKLYKNCPKAMKRLVADCVKKVKEERPLFP
QILSSIELLQHSLPKINRSASEPSLHRAAHTEDINACTLT
TSPRLPVFYSFLPFFFFFFSFCTPSTF

FIGURE 15





FIGURE 16







FIGURE 17




G12V-Ras

G12V-Ras + FRNK



Raf-caax



Raf-caax + FRNK





FIGURE 18

A




B

bFGF



C

G12V-Ras



D

Raf-caax




+ empty
vector

E



F



G



+ FRNK

FIGURE 19




FIGURE 20A




FIGURE 20B

aaa ctc aca gat cct tct aag aca agc aac act atc cgt qtt ttc ttg 315

Lys	Leu	Thr	Asp	Pro	Ser	Lys	Thr	Ser	Asn	Thr	Ile	Arg	Val	Phe	Leu	
			50						55				60			
ccg	aac	aag	caa	aga	aca	gtg	gtc	aat	gtg	cga	aat	gga	atg	agc	ttg	363
Pro	Asn	Lys	Gln	Arg	Thr	Val	Val	Asn	Val	Arg	Asn	Gly	Met	Ser	Leu	
		65					70					75				
cat	gac	tgc	ctt	atg	aaa	gca	ctc	aag	gtg	agg	ggc	ctg	caa	cca	gag	411
His	Asp	Cys	Leu	Met	Lys	Ala	Leu	Lys	Val	Arg	Gly	Leu	Gln	Pro	Glu	
	80					85					90					
tgc	tgt	gca	gtg	ttc	aga	ctt	ctc	cac	gaa	cac	aaa	ggg	aaa	aaa	gca	459
Cys	Cys	Ala	Val	Phe	Arg	Leu	Leu	His	Glu	His	Lys	Gly	Lys	Lys	Ala	
95					100					105					110	
cgc	tta	gat	tgg	aat	act	gat	gct	gcg	tct	ttg	att	gga	gaa	gaa	ctt	507
Arg	Leu	Asp	Trp	Asn	Thr	Asp	Ala	Ala	Ser	Leu	Ile	Gly	Glu	Glu	Leu	
			115						120					125		
caa	gta	gat	ttc	ctg	gat	cat	gtt	ccc	ctc	aca	aca	cac	aac	ttt	gct	555
Gln	Val	Asp	Phe	Leu	Asp	His	Val	Pro	Leu	Thr	Thr	His	Asn	Phe	Ala	
			130					135					140			
cgg	aag	acg	ttc	ctg	aag	ctt	gcc	ttc	tgt	gac	atc	tgt	cag	aaa	ttc	603
Arg	Lys	Thr	Phe	Leu	Lys	Leu	Ala	Phe	Cys	Asp	Ile	Cys	Gln	Lys	Phe	
		145					150					155				
ctg	ctc	aat	gga	ttt	cga	tgt	cag	act	tgt	ggc	tac	aaa	ttt	cat	gag	651
Leu	Leu	Asn	Gly	Phe	Arg	Cys	Gln	Thr	Cys	Gly	Tyr	Lys	Phe	His	Glu	
	160					165					170					
cac	tgt	agc	acc	aaa	gta	cct	act	atg	tgt	gtg	gac	tgg	agt	aac	atc	699
His	Cys	Ser	Thr	Lys	Val	Pro	Thr	Met	Cys	Val	Asp	Trp	Ser	Asn	Ile	
175					180					185					190	
aga	caa	ctc	tta	ttg	ttt	cca	aat	tcc	act	att	ggg	gat	agt	gga	gtc	747
Arg	Gln	Leu	Leu	Leu	Phe	Pro	Asn	Ser	Thr	Ile	Gly	Asp	Ser	Gly	Val	
			195						200					205		
cca	gca	cta	cct	tct	ttg	act	atg	cgt	cgt	atg	cga	gag	tct	gtt	tcc	795
Pro	Ala	Leu	Pro	Ser	Leu	Thr	Met	Arg	Arg	Met	Arg	Glu	Ser	Val	Ser	
			210					215					220			
agg	atg	cct	gtt	agt	tct	cag	cac	aga	tat	tct	aca	cct	cac	gcc	ttc	843
Arg	Met	Pro	Val	Ser	Ser	Gln	His	Arg	Tyr	Ser	Thr	Pro	His	Ala	Phe	
		225					230					235				
acc	ttt	aac	acc	tcc	agt	ccc	tca	tct	gaa	ggg	tcc	ctc	tcc	cag	agg	891
Thr	Phe	Asn	Thr	Ser	Ser	Pro	Ser	Ser	Glu	Gly	Ser	Leu	Ser	Gln	Arg	
	240					245					250					

cag	agg	tcg	aca	tcc	aca	cct	aat	gtc	cac	atg	gtc	agc	acc	acg	ctg	939
Gln	Arg	Ser	Thr	Ser	Thr	Pro	Asn	Val	His	Met	Val	Ser	Thr	Thr	Leu	
255					260					265					270	
cct	gtg	gac	agc	agg	atg	att	gag	gat	gca	att	cga	agt	cac	agc	gaa	987
Pro	Val	Asp	Ser	Arg	Met	Ile	Glu	Asp	Ala	Ile	Arg	Ser	His	Ser	Glu	
				275					280					285		
tca	gcc	tca	cct	tca	gcc	ctg	tcc	agt	agc	ccc	aac	aat	ctg	agc	cca	1035
Ser	Ala	Ser	Pro	Ser	Ala	Leu	Ser	Ser	Ser	Pro	Asn	Asn	Leu	Ser	Pro	
			290					295					300			
aca	ggc	tgg	tca	cag	ccg	aaa	acc	ccc	gtg	cca	gca	caa	aga	gag	cgg	1083
Thr	Gly	Trp	Ser	Gln	Pro	Lys	Thr	Pro	Val	Pro	Ala	Gln	Arg	Glu	Arg	
		305				310						315				
gca	cca	gta	tct	ggg	acc	cag	gag	aaa	aac	aaa	att	agg	cct	cgt	gga	1131
Ala	Pro	Val	Ser	Gly	Thr	Gln	Glu	Lys	Asn	Lys	Ile	Arg	Pro	Arg	Gly	
	320					325				330						
cag	aga	gat	tca	agc	tat	tat	tgg	gaa	ata	gaa	gcc	agt	gaa	gtg	atg	1179
Gln	Arg	Asp	Ser	Ser	Tyr	Tyr	Trp	Glu	Ile	Glu	Ala	Ser	Glu	Val	Met	
335					340					345					350	
ctg	tcc	act	cgg	att	ggg	tca	ggc	tct	ttt	gga	act	gtt	tat	aag	ggt	1227
Leu	Ser	Thr	Arg	Ile	Gly	Ser	Gly	Ser	Phe	Gly	Thr	Val	Tyr	Lys	Gly	
				355					360					365		
aaa	tgg	cac	gga	gat	gtt	gca	gta	aag	atc	cta	aag	gtt	gtc	gac	cca	1275
Lys	Trp	His	Gly	Asp	Val	Ala	Val	Lys	Ile	Leu	Lys	Val	Val	Asp	Pro	
			370					375					380			
acc	cca	gag	caa	ttc	cag	gcc	ttc	agg	aat	gag	gtg	gct	gtt	ctg	cgc	1323
Thr	Pro	Glu	Gln	Phe	Gln	Ala	Phe	Arg	Asn	Glu	Val	Ala	Val	Leu	Arg	
		385					390					395				
aaa	aca	cgg	cat	gtg	aac	att	ctg	ctt	ttc	atg	ggg	tac	atg	aca	aag	1371
Lys	Thr	Arg	His	Val	Asn	Ile	Leu	Leu	Phe	Met	Gly	Tyr	Met	Thr	Lys	
	400					405					410					
gac	aac	ctg	gca	att	gtg	acc	cag	tgg	tgc	gag	ggc	agc	agc	ctc	tac	1419
Asp	Asn	Leu	Ala	Ile	Val	Thr	Gln	Trp	Cys	Glu	Gly	Ser	Ser	Leu	Tyr	
415					420					425					430	
aaa	cac	ctg	cat	gtc	cag	gag	acc	aag	ttt	cag	atg	ttc	cag	cta	att	1467
Lys	His	Leu	His	Val	Gln	Glu	Thr	Lys	Phe	Gln	Met	Phe	Gln	Leu	Ile	
				435					440					445		
gac	att	gcc	cgg	cag	acg	gct	cag	gga	atg	gac	tat	ttg	cat	gca	aag	1515
Asp	Ile	Ala	Arg	Gln	Thr	Ala	Gln	Gly	Met	Asp	Tyr	Leu	His	Ala	Lys	
			450					455					460			

aac atc atc cat aga gac atg aaa tcc aac aat ata ttt ctc cat gaa 1563
 Asn Ile Ile His Arg Asp Met Lys Ser Asn Asn Ile Phe Leu His Glu
 465 470 475

ggc tta aca gtg aaa att gga gat ttt ggt ttg gca aca gta aag tca 1611
 Gly Leu Thr Val Lys Ile Gly Asp Phe Gly Leu Ala Thr Val Lys Ser
 480 485 490

cgc tgg agt ggt tct cag cag gtt gaa caa cct act ggc tct gtc ctc 1659
 Arg Trp Ser Gly Ser Gln Gln Val Glu Gln Pro Thr Gly Ser Val Leu
 495 500 505 510

tgg atg gcc cca gag gtg atc cga atg cag gat aac aac cca ttc agt 1707
 Trp Met Ala Pro Glu Val Ile Arg Met Gln Asp Asn Asn Pro Phe Ser
 515 520 525

ttc cag tcg gat gtc tac tcc tat ggc atc gta ttg tat gaa ctg atg 1755
 Phe Gln Ser Asp Val Tyr Ser Tyr Gly Ile Val Leu Tyr Glu Leu Met
 530 535 540

acg ggg gag ctt cct tat tct cac atc aac aac cga gat cag atc atc 1803
 Thr Gly Glu Leu Pro Tyr Ser His Ile Asn Asn Arg Asp Gln Ile Ile
 545 550 555

ttc atg gtg ggc cga gga tat gcc tcc cca gat ctt agt aag cta tat 1851
 Phe Met Val Gly Arg Gly Tyr Ala Ser Pro Asp Leu Ser Lys Leu Tyr
 560 565 570

aag aac tgc ccc aaa gca atg aag agg ctg gta gct gac tgt gtg aag 1899
 Lys Asn Cys Pro Lys Ala Met Lys Arg Leu Val Ala Asp Cys Val Lys
 575 580 585 590

aaa gta aag gaa gag agg cct ctt ttt ccc cag atc ctg tct tcc att 1947
 Lys Val Lys Glu Glu Arg Pro Leu Phe Pro Gln Ile Leu Ser Ser Ile
 595 600 605

gag ctg ctc caa cac tct cta ccg aag atc aac cgg agc gct tcc gag 1995
 Glu Leu Leu Gln His Ser Leu Pro Lys Ile Asn Arg Ser Ala Ser Glu
 610 615 620

cca tcc ttg cat cgg gca gcc cac act gag gat atc aat gct tgc acg 2043
 Pro Ser Leu His Arg Ala Ala His Thr Glu Asp Ile Asn Ala Cys Thr
 625 630 635

ctg acc acg tcc ccg agg ctg cct gtc ttc tagttgactt tgcacctgctc 2093
 Leu Thr Thr Ser Pro Arg Leu Pro Val Phe
 640 645

ttcaggctgc caggggagga ggagaagcca gcaggcacca cttttctgct ccctttctcc 2153

```
<210> 2
<211> 648
<212> PRT
<213> Homo sapiens
```

Met Glu His Ile Gln Gly Ala Trp Lys Thr Ile Ser Asn Gly Phe Gly
1 5 10 15

Phe Lys Asp Ala Val Phe Asp Gly Ser Ser Cys Ile Ser Pro Thr Ile
20 25 30

Val Gln Gln Phe Gly Tyr Gln Arg Arg Ala Ser Asp Asp Gly Lys Leu
35 40 45

Thr Asp Pro Ser Lys Thr Ser Asn Thr Ile Arg Val Phe Leu Pro Asn
50 55 60

Lys Gln Arg Thr Val Val Asn Val Arg Asn Gly Met Ser Leu His Asp
65 70 75 80

Cys Leu Met Lys Ala Leu Lys Val Arg Gly Leu Gln Pro Glu Cys Cys

6/20

85					90					95					
Ala	Val	Phe	Arg	Leu	Leu	His	Glu	His	Lys	Gly	Lys	Lys	Ala	Arg	Leu
			100					105					110		
Asp	Trp	Asn	Thr	Asp	Ala	Ala	Ser	Leu	Ile	Gly	Glu	Glu	Leu	Gln	Val
		115					120					125			
Asp	Phe	Leu	Asp	His	Val	Pro	Leu	Thr	Thr	His	Asn	Phe	Ala	Arg	Lys
	130					135					140				
Thr	Phe	Leu	Lys	Leu	Ala	Phe	Cys	Asp	Ile	Cys	Gln	Lys	Phe	Leu	Leu
145					150					155					160
Asn	Gly	Phe	Arg	Cys	Gln	Thr	Cys	Gly	Tyr	Lys	Phe	His	Glu	His	Cys
				165					170					175	
Ser	Thr	Lys	Val	Pro	Thr	Met	Cys	Val	Asp	Trp	Ser	Asn	Ile	Arg	Gln
			180						185				190		
Leu	Leu	Leu	Phe	Pro	Asn	Ser	Thr	Ile	Gly	Asp	Ser	Gly	Val	Pro	Ala
		195					200					205			
Leu	Pro	Ser	Leu	Thr	Met	Arg	Arg	Met	Arg	Glu	Ser	Val	Ser	Arg	Met
	210					215					220				
Pro	Val	Ser	Ser	Gln	His	Arg	Tyr	Ser	Thr	Pro	His	Ala	Phe	Thr	Phe
225						230					235				240
Asn	Thr	Ser	Ser	Pro	Ser	Ser	Glu	Gly	Ser	Leu	Ser	Gln	Arg	Gln	Arg
				245					250					255	
Ser	Thr	Ser	Thr	Pro	Asn	Val	His	Met	Val	Ser	Thr	Thr	Leu	Pro	Val
			260					265					270		
Asp	Ser	Arg	Met	Ile	Glu	Asp	Ala	Ile	Arg	Ser	His	Ser	Glu	Ser	Ala
		275					280					285			
Ser	Pro	Ser	Ala	Leu	Ser	Ser	Ser	Pro	Asn	Asn	Leu	Ser	Pro	Thr	Gly
	290					295					300				
Trp	Ser	Gln	Pro	Lys	Thr	Pro	Val	Pro	Ala	Gln	Arg	Glu	Arg	Ala	Pro
305						310					315				320
Val	Ser	Gly	Thr	Gln	Glu	Lys	Asn	Lys	Ile	Arg	Pro	Arg	Gly	Gln	Arg
				325					330					335	
Asp	Ser	Ser	Tyr	Tyr	Trp	Glu	Ile	Glu	Ala	Ser	Glu	Val	Met	Leu	Ser
			340					345				350			
Thr	Arg	Ile	Gly	Ser	Gly	Ser	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Lys	Trp

DOT PLOT ANALYSIS

355	360	365
His Gly Asp Val Ala Val Lys Ile Leu Lys Val Val Asp Pro Thr Pro		
370	375	380
Glu Gln Phe Gln Ala Phe Arg Asn Glu Val Ala Val Leu Arg Lys Thr		
385	390	395
Arg His Val Asn Ile Leu Leu Phe Met Gly Tyr Met Thr Lys Asp Asn		
	405	410
		415
Leu Ala Ile Val Thr Gln Trp Cys Glu Gly Ser Ser Leu Tyr Lys His		
	420	425
		430
Leu His Val Gln Glu Thr Lys Phe Gln Met Phe Gln Leu Ile Asp Ile		
	435	440
		445
Ala Arg Gln Thr Ala Gln Gly Met Asp Tyr Leu His Ala Lys Asn Ile		
	450	455
		460
Ile His Arg Asp Met Lys Ser Asn Asn Ile Phe Leu His Glu Gly Leu		
465	470	475
		480
Thr Val Lys Ile Gly Asp Phe Gly Leu Ala Thr Val Lys Ser Arg Trp		
	485	490
		495
Ser Gly Ser Gln Gln Val Glu Gln Pro Thr Gly Ser Val Leu Trp Met		
	500	505
		510
Ala Pro Glu Val Ile Arg Met Gln Asp Asn Asn Pro Phe Ser Phe Gln		
	515	520
		525
Ser Asp Val Tyr Ser Tyr Gly Ile Val Leu Tyr Glu Leu Met Thr Gly		
	530	535
		540
Glu Leu Pro Tyr Ser His Ile Asn Asn Arg Asp Gln Ile Ile Phe Met		
545	550	555
		560
Val Gly Arg Gly Tyr Ala Ser Pro Asp Leu Ser Lys Leu Tyr Lys Asn		
	565	570
		575
Cys Pro Lys Ala Met Lys Arg Leu Val Ala Asp Cys Val Lys Lys Val		
	580	585
		590
Lys Glu Glu Arg Pro Leu Phe Pro Gln Ile Leu Ser Ser Ile Glu Leu		
	595	600
		605
Leu Gln His Ser Leu Pro Lys Ile Asn Arg Ser Ala Ser Glu Pro Ser		
	610	615
		620
Leu His Arg Ala Ala His Thr Glu Asp Ile Asn Ala Cys Thr Leu Thr		

001130-3064950

8/20

625

630

635

640

Thr Ser Pro Arg Leu Pro Val Phe
645

<210> 3

<211> 570

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(567)

<400> 3

atg acg gaa tat aag ctg gtg gtg gtg ggc gcc ggc ggt gtg ggc aag	48
Met Thr Glu Tyr Lys Leu Val Val Val Gly Ala Gly Gly Val Gly Lys	
1 5 10 15	
agt gcg ctg acc atc cag ctg atc cag aac cat ttt gtg gac gaa tac	96
Ser Ala Leu Thr Ile Gln Leu Ile Gln Asn His Phe Val Asp Glu Tyr	
20 25 30	
gac ccc act ata gag gat tcc tac cgg aag cag gtg gtc att gat ggg	144
Asp Pro Thr Ile Glu Asp Ser Tyr Arg Lys Gln Val Val Ile Asp Gly	
35 40 45	
gag acg tgc ctg ttg gac atc ctg gat acc gcc ggc cag gag gag tac	192
Glu Thr Cys Leu Leu Asp Ile Leu Asp Thr Ala Gly Gln Glu Glu Tyr	
50 55 60	
agc gcc atg cgg gac cag tac atg cgc acc ggg gag ggc ttc ctg tgt	240
Ser Ala Met Arg Asp Gln Tyr Met Arg Thr Gly Glu Gly Phe Leu Cys	
65 70 75 80	
gtg ttt gcc atc aac aac acc aag tct ttt gag gac atc cac cag tac	288
Val Phe Ala Ile Asn Asn Thr Lys Ser Phe Glu Asp Ile His Gln Tyr	
85 90 95	
agg gag cag atc aaa cgg gtg aag gac tcg gat gac gtg ccc atg gtg	336
Arg Glu Gln Ile Lys Arg Val Lys Asp Ser Asp Asp Val Pro Met Val	
100 105 110	
ctg gtg ggg aac aag tgt gac ctg gct gca cgc act gtg gaa tct cgg	384
Leu Val Gly Asn Lys Cys Asp Leu Ala Ala Arg Thr Val Glu Ser Arg	
115 120 125	
cag gct cag gac ctc gcc cga agc tac ggc atc ccc tac atc gag acc	432
Gln Ala Gln Asp Leu Ala Arg Ser Tyr Gly Ile Pro Tyr Ile Glu Thr	
130 135 140	

001100-001100

tcg gcc aag acc cgg cag gga gtg gag gat gcc ttc tac acg ttg gtg 480
 Ser Ala Lys Thr Arg Gln Gly Val Glu Asp Ala Phe Tyr Thr Leu Val
 145 150 155 160

cgt gag atc cgg cag cac aag ctg cgg aag ctg aac cct cct gat gag 528
 Arg Glu Ile Arg Gln His Lys Leu Arg Lys Leu Asn Pro Pro Asp Glu
 165 170 175

agt ggc ccc ggc tgc atg agc tgc aag tgt gtg ctc tcc tga 570
 Ser Gly Pro Gly Cys Met Ser Cys Lys Cys Val Leu Ser
 180 185

<210> 4

<211> 189

<212> PRT

<213> Homo sapiens

<400> 4

Met Thr Glu Tyr Lys Leu Val Val Val Gly Ala Gly Gly Val Gly Lys
 1 5 10 15

Ser Ala Leu Thr Ile Gln Leu Ile Gln Asn His Phe Val Asp Glu Tyr
 20 25 30

Asp Pro Thr Ile Glu Asp Ser Tyr Arg Lys Gln Val Val Ile Asp Gly
 35 40 45

Glu Thr Cys Leu Leu Asp Ile Leu Asp Thr Ala Gly Gln Glu Glu Tyr
 50 55 60

Ser Ala Met Arg Asp Gln Tyr Met Arg Thr Gly Glu Gly Phe Leu Cys
 65 70 75 80

Val Phe Ala Ile Asn Asn Thr Lys Ser Phe Glu Asp Ile His Gln Tyr
 85 90 95

Arg Glu Gln Ile Lys Arg Val Lys Asp Ser Asp Asp Val Pro Met Val
 100 105 110

Leu Val Gly Asn Lys Cys Asp Leu Ala Ala Arg Thr Val Glu Ser Arg
 115 120 125

Gln Ala Gln Asp Leu Ala Arg Ser Tyr Gly Ile Pro Tyr Ile Glu Thr
 130 135 140

Ser Ala Lys Thr Arg Gln Gly Val Glu Asp Ala Phe Tyr Thr Leu Val
 145 150 155 160

Arg Glu Ile Arg Gln His Lys Leu Arg Lys Leu Asn Pro Pro Asp Glu

001130-206000

165

170

175

Ser Gly Pro Gly Cys Met Ser Cys Lys Cys Val Leu Ser
 180 185

<210> 5

<211> 6453

<212> DNA

<213> Homo sapiens

<220>

<221> prim_transcript

<222> (1664)..(3744)

<220>

<221> intron

<222> (1775)..(2041)

<220>

<221> intron

<222> (2221)..(2373)

<220>

<221> intron

<222> (2534)..(3230)

<400> 5

ggatcccagc ctttccccag cccgtagccc cgggacctcc gcggtgggcg gcgccgcgct 60
 gccggcgagc ggagggcctc tgggtgcaccg gcaccgctga gtcgggttct ctcgccggcc 120
 tgttcccggg agagcccggg gccctgctcg gagatgccgc cccggggccc cagacaccgg 180
 ctccctggcc ttcctcgagc aaccccagagc tcggctccgg tctccagcca agcccaacct 240
 cgagaggccg cggccctact ggctccgcct cccgcgttgc tcccggaagc cccgcccgac 300
 cgcggtcctt gacagacggg ccgctcagcc aaccgggggtg gggcggggcc cgatggcgcg 360
 cagccaatgg taggccgcgc ctggcagacg gacggggcgcg gggcggggcg tgcgcaggcc 420
 cgcccagatc tccgccgccc gtgccctgcg cccgcaacct gagccgcacc cgccgcggac 480
 ggagcccatg cgcggggcga accgcgcgcc cccgcccccg ccccgccccg gcctcggccc 540
 cggccctggc cccgggggca gtcgcgcctg tgaacggtga gtgcgggcag ggatcggccg 600
 ggccgcgcgc cctcctcgcc cccaggcggc agcaatacgc gcggcgcggg ccgggggcgc 660
 ggggcccggc ggcgtaagcg gcggcggcgg cggcggggtg gtggggccgg gcggggcccc 720

cgggcacagg tgagcgggcg tcgggggctg cggcgggcg gggccccttc ctccctgggg 780
 cctgcgggaa tccgggcccc acccgtggcc tcgcgctggg cacggtcccc acgccggcgt 840
 acccgggagc ctccgggccc ggcacctcac acccgggggc gtctgggagg aggcggccgc 900
 ggccacggca cggccgggca cccccgattc agcatcacag gtcgcggacc aggccggggg 960
 cctcagcccc agtgcccttt ccctctccgg gtctcccgcg ccgcttctcg gccccttcct 1020
 gtgcgtcagt ccctgcttcc caggagctcc tctgtcttct ccagctttct gtggctgaaa 1080
 gatgcccccg gttcccgcgc ggggggtgcg ggcgctgccc gggctgccc tcccctcggc 1140
 ggcgcctagt acgcagtagg cgctcagcaa atactgtcg gaggcaccag cgccgcgggg 1200
 cctgcaggct ggcactagcc tgcccgggca cgcgctggcg cgctccgccg tggccagacc 1260
 tgttctggag gacggtaacc tcagccctcg ggcgcctccc tttagccttt ctgccgaccc 1320
 agcagcttct aatttggtg cgtgggtgag agcgtcagc tgtcagccct gcctttgagg 1380
 gctgggtccc ttttcccatc actgggtcat taagagcaag tgggggcgag gcgacagccc 1440
 tcccgcacgc tgggttgag ctgcacaggt aggcacgctg cagtccttgc tgcctggcgt 1500
 tggggcccag ggaccgctgt gggtttgccc ttcagatggc cctgccagca gctgccctgt 1560
 ggggcctggg gctgggcctg ggcctggctg agcagggccc tccttggcag gtggggcagg 1620
 agaccctgta ggaggacccc gggccgcagg cccctgagga gcgatgacgg aatataagct 1680
 ggtggtggtg ggcgccggcg gtgtgggcaa gagtgcgctg accatccagc tgatccagaa 1740
 ccattttgtg gacgaatacg accccactat agaggtgagc ctagcgccgc cgtccagggt 1800
 ccagcagctg ctgcgggcca gcccaggaca cagccaggat agggctggct gcagcccctg 1860
 gtcccctgca tgggtgctgt gccctgtctc ctgcttcctc tagaggaggg gagtcctcg 1920
 tctcagcacc ccaggagagg agggggcatg aggggcatga gaggtaccag ggagaggctg 1980
 gctgtgtgaa cccccccac ggaaggctct gagggggtcc ctgagccctg tcctcctgca 2040
 ggattcctac cggaagcagg tggtcattga tggggagacg tgctgttgg acatcctgga 2100
 taccgccggc caggaggagt acagcgccat gcgggaccag tacatgcgca ccggggaggg 2160
 cttcctgtgt gtgtttgcca tcaacaacac caagtctttt gaggacatcc accagtacag 2220

gtgaaccccg tgaggctggc ccgggagccc acgccgcaca ggtggggcca ggccggctgc 2280
 gtccaggcag gggcctcctg tcctctctgc gcatgtcctg gatgccgctg cgctgcagc 2340
 ccccgtagcc agctctcgct ttccacctct cagggagcag atcaaacggg tgaaggactc 2400
 ggatgacgtg cccatggtgc tgggtgggaa caagtgtgac ctggctgcac gcaactgtgga 2460
 atctcggcag gctcaggacc tcgcccgaag ctacggcatc ccctacatcg agacctcggc 2520
 caagaccccg caggtgaggc agctctccac ccacagcta gccagggacc cgcgccgccc 2580
 cgcgccagcc agggagcagc actcactgac cctctccctt gacacagggc agccgctctg 2640
 gctctagctc cagctccggg accctctggg acccccggg acccatgtga ccagcggcc 2700
 cctcgcactg taggtctccc gggacggcag ggcagtgagg gaggcgaggg ccggggctctg 2760
 ggctcacgcc ctgcagtcc tggccgacac agctccgggg aaggcgaggg tccttgggga 2820
 gagctgccct gagccaggcc ggagcgggtga ccctggggcc cggcccctct tgtcccaga 2880
 gtgtcccacg ggcacctgtt ggttctgagt cttagtgggg ctactgggga cacgggccgt 2940
 agctgagtcg agagctgggt gcagggtggt caaacctgg ccagacctgg agttcaggag 3000
 ggccccgggc caccctgacc tttagggggc tgctgtagca tgatgcgggt ggccctgggc 3060
 acttcgagat ggccagagtc cagcttcccg tgtgtgtggt gggcctgggg aagtggctgg 3120
 tggagtcggg agcttcgggc caggcaaggc ttgatccac agcagggagc ccctcaccca 3180
 ggcaggcggc cacaggccgg tcctcctga tccatccct cctttcccag ggagtggagg 3240
 atgccttcta cacgttggtg cgtgagatcc ggcagcaca gctgcggaag ctgaaccctc 3300
 ctgatgagag tggccccggc tgcattgagc gcaagtgtgt gctctcctga cgcagggtgag 3360
 ggggactccc agggcgcccg ccacgcccac cggatgacct cggctccccg ccctgcccgg 3420
 tctcctggcc tgcggtcagc agcctccctt gtgccccgcc cagcacaagc tcaggacatg 3480
 gaggtgccgg atgcaggaag gaggtgcaga cggaaggagg aggaaggaag gacggaagca 3540
 aggaaggaag gaagggtctg tggagcccag tcaccccggg accgtggggc gaggtgactg 3600
 cagaccctcc cagggaggct gtgcacagac tgtcttgaac atcccaaag ccaccggaac 3660
 cccagccctt agctcccctc ccaggcctct gtgggccctt gtcgggcaca gatgggatca 3720
 cagtaaatta ttgatgggtc ttgatcttgg ttttcggctg aggggtgggac acgggtgcgcg 3780

tgtggcctgg catgaggtat gtcggaacct caggcctgtc cagccctggg ctctccatag 3840
 cctttgggag ggggaggttg ggagaggccg gtcaggggtc tgggctgtgg tgctctctcc 3900
 tccgcctgc cccagtgtcc acggcttctg gcagagagct ctggacaagc aggcagatca 3960
 taaggacaga gagcttactg tgcttctacc aactaggagg gcgtcctggt cctccagagg 4020
 gaggtggttt caggggttgg ggatctgtgc cggtaggctct ggtctctgct gggagccttc 4080
 ttggcgggtga gaggcatac ctttctgac ttgctcccag cgtgaaatgc acctgccaag 4140
 aatggcagac atagggaccc cgcctcctgg gccttcacat gccagtttt cttcggctct 4200
 gtggcctgaa gcggtctgtg gaccttgga gtagggctcc agcaccgact ggcctcaggc 4260
 ctctgcctca ttggtggtcg ggtagcggcc agtagggcgt gggagcctgg ccatccctgc 4320
 ctctggaggt ggacgaggtt ggcagctggt ccgtctgctc ctgccccact ccccccgcc 4380
 cctgcctca cctaccctt gccccacgcc tgcctcatgg ctggttgctc ttggagcctg 4440
 gtagtgtcac tggctcagcc ttgctgggta tacacaggct ctgccacca ctctgctcca 4500
 aggggcttgc cctgccttgg gccaaagtct aggtctggcc acagccacag acagctcagt 4560
 ccctgtgtg gtcaccttgg cttctgctgg gggccacag cgccttgggt gcccctccc 4620
 tcccagggcc cgggttgagg ctgggccagg cctctggga cggggacttg tgccctgtca 4680
 gggttcccta tccctgaggt tgggggagag ctagcagggc atgccgctgg ctggccaggg 4740
 ctgcaggac actccccctt ttgtccagg aataccacac tcgcccttct ctccagcgaa 4800
 caccacactc gcccttctct ccaggggacg ccacactccc cttctgtcc aggggacgcc 4860
 aactcccc ttctctccag gggacgccac actcgccctt ctctccagg gacgccacac 4920
 tcgcccttct ctccagggga cgccacactc gcccttctgt ccaggggacg ccacactcgc 4980
 cttctctcc aggggacgcc aactcgccc ttctctccag gggacgccac actccccctt 5040
 ctgtccagg gacgccacac tcccccttct ctccagggga cgccacactc ccccttctct 5100
 ccaggggacg ccacactcgc cttctctcc aggggacgcc aactcccc ttctgtccag 5160
 gggacgccac actcgccctt ctctccagg gacgccacac tcgcccttct ctccagggga 5220
 cgccacactc ccccttctct ccaggggacg ccacactccc cttctctcc aggggacgcc 5280

acactccccc ttctgtccag gggacgccac actcgccctt ctctccaggg gacgccacac 5340
 tcccccttct ctccagggga cgccacactc ccccttctct ccaggggacg ccacactccc 5400
 ccttctgtcc aggggacgcc acactcgccc ttctctccag gggacgccac actcgccctt 5460
 ctctccaggg gacgccacac tcgcccttct ctccagggga cgccacactt gcccttctgt 5520
 ccagggaatg ccacactccc ccttctcccc agcagcctcc gactgaccag cttccccatc 5580
 gatagacttc ccgaggccag gagccctcta gggctgccgg gtgccaccct ggctccttcc 5640
 acaccgtgct ggtcactgcc tgctgggggc gtcagatgca ggtgaccctg tgcaggaggt 5700
 atctctggac ctgcctcttg gtcattacgg ggctgggcag ggctggtat cagggccccc 5760
 ctggggttgc agggctgggc ctgtgctgtg gtcctggggg gtccaggaca gacgtggagg 5820
 ggtcagggcc cagcaccctt gtcctatgct gaactgtggg aagcatccag gtccctgggt 5880
 ggcttcaaca ggagttccag cacgggaacc actggacaac ctgggggtgtg tcctgatctg 5940
 gggacaggcc agccacaccc cgagtcctag ggactccaga gagcagccca ctgccctggg 6000
 ctccacggaa gccccctcat gccgctaggc cttggcctcg gggacagccc agctaggcca 6060
 gtgtgtggca ggaccaggcc cccatgtggg agctgacccc ttgggattct ggagctgtgc 6120
 tgatgggcag gggagagcca gtcctcccc ttgaggagg gtcttgatgc ctggggttac 6180
 ccgcagaggc ctgggtgccg ggacgctccc cggtttggt gaaaggaaag cagatgtggg 6240
 cagcttctcc actgagccca tctggtcttc ccggggctgg gcccataga tctgggtccc 6300
 tgtgtggccc ccctggtctg atgccgagga taccctgca aactgccaat ccagaggac 6360
 aagactggga agtcctgca gggagagccc atccccgcac cctgaccac aagagggact 6420
 cctgctgccc accaggcatc cctccagga tcc 6453

<210> 6

<211> 2004

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: fusion protein

<220>

<221> CDS

<222> (1)..(2004)

<400> 6

atg gag cac ata cag gga gct tgg aag acg atc agc aat ggt ttt gga	48
Met Glu His Ile Gln Gly Ala Trp Lys Thr Ile Ser Asn Gly Phe Gly	
1 5 10 15	
ttc aaa gat gcc gtg ttt gat ggc tcc agc tgc atc tct cct aca ata	96
Phe Lys Asp Ala Val Phe Asp Gly Ser Ser Cys Ile Ser Pro Thr Ile	
20 25 30	
gtt cag cag ttt ggc tat cag cgc cgg gca tca gat gat ggc aaa ctc	144
Val Gln Gln Phe Gly Tyr Gln Arg Arg Ala Ser Asp Asp Gly Lys Leu	
35 40 45	
aca gat cct tct aag aca agc aac act atc cgt gtt ttc ttg ccg aac	192
Thr Asp Pro Ser Lys Thr Ser Asn Thr Ile Arg Val Phe Leu Pro Asn	
50 55 60	
aag caa aga aca gtg gtc aat gtg cga aat gga atg agc ttg cat gac	240
Lys Gln Arg Thr Val Val Asn Val Arg Asn Gly Met Ser Leu His Asp	
65 70 75 80	
tgc ctt atg aaa gca ctc aag gtg agg ggc ctg caa cca gag tgc tgt	288
Cys Leu Met Lys Ala Leu Lys Val Arg Gly Leu Gln Pro Glu Cys Cys	
85 90 95	
gca gtg ttc aga ctt ctc cac gaa cac aaa ggt aaa aaa gca cgc tta	336
Ala Val Phe Arg Leu Leu His Glu His Lys Gly Lys Lys Ala Arg Leu	
100 105 110	
gat tgg aat act gat gct gcg tct ttg att gga gaa gaa ctt caa gta	384
Asp Trp Asn Thr Asp Ala Ala Ser Leu Ile Gly Glu Glu Leu Gln Val	
115 120 125	
gat ttc ctg gat cat gtt ccc ctc aca aca cac aac ttt gct cgg aag	432
Asp Phe Leu Asp His Val Pro Leu Thr Thr His Asn Phe Ala Arg Lys	
130 135 140	
acg ttc ctg aag ctt gcc ttc tgt gac atc tgt cag aaa ttc ctg ctc	480
Thr Phe Leu Lys Leu Ala Phe Cys Asp Ile Cys Gln Lys Phe Leu Leu	
145 150 155 160	
aat gga ttt cga tgt cag act tgt ggc tac aaa ttt cat gag cac tgt	528
Asn Gly Phe Arg Cys Gln Thr Cys Gly Tyr Lys Phe His Glu His Cys	
165 170 175	
agc acc aaa gta cct act atg tgt gtg gac tgg agt aac atc aga caa	576
Ser Thr Lys Val Pro Thr Met Cys Val Asp Trp Ser Asn Ile Arg Gln	
180 185 190	

ctc Leu	tta Leu	ttg Leu	ttt Phe	cca Pro	aat Asn	tcc Ser	act Thr	att Ile	ggt Gly	gat Asp	agt Ser	gga Gly	gtc Val	cca Pro	gca Ala	624
195						200						205				
cta Leu	cct Pro	tct Ser	ttg Leu	act Thr	atg Met	cgt Arg	cgt Arg	atg Met	cga Arg	gag Glu	tct Ser	gtt Val	tcc Ser	agg Arg	atg Met	672
210						215						220				
cct Pro	gtt Val	agt Ser	tct Ser	cag Gln	cac His	aga Arg	tat Tyr	tct Ser	aca Thr	cct Pro	cac His	gcc Ala	ttc Phe	acc Thr	ttt Phe	720
225						230						235			240	
aac Asn	acc Thr	tcc Ser	agt Ser	ccc Pro	tca Ser	tct Ser	gaa Glu	ggt Gly	tcc Ser	ctc Leu	tcc Ser	cag Gln	agg Arg	cag Gln	agg Arg	768
			245						250						255	
tcg Ser	aca Thr	tcc Ser	aca Thr	cct Pro	aat Asn	gtc Val	cac His	atg Met	gtc Val	agc Ser	acc Thr	acg Thr	ctg Leu	cct Pro	gtg Val	816
			260						265						270	
gac Asp	agc Ser	agg Arg	atg Met	att Ile	gag Glu	gat Asp	gca Ala	att Ile	cga Arg	agt Ser	cac His	agc Ser	gaa Glu	tca Ser	gcc Ala	864
			275						280						285	
tca Ser	cct Pro	tca Ser	gcc Ala	ctg Leu	tcc Ser	agt Ser	agc Ser	ccc Pro	aac Asn	aat Asn	ctg Leu	agc Ser	cca Pro	aca Thr	ggc Gly	912
290						295						300				
tgg Trp	tca Ser	cag Gln	ccg Pro	aaa Lys	acc Thr	ccc Pro	gtg Val	cca Pro	gca Ala	caa Gln	aga Arg	gag Glu	cgg Arg	gca Ala	cca Pro	960
305						310						315			320	
gta Val	tct Ser	ggg Gly	acc Thr	cag Gln	gag Glu	aaa Lys	aac Asn	aaa Lys	att Ile	agg Arg	cct Pro	cgt Arg	gga Gly	cag Gln	aga Arg	1008
			325						330						335	
gat Asp	tca Ser	agc Ser	tat Tyr	tat Tyr	tgg Trp	gaa Glu	ata Ile	gaa Glu	gcc Ala	agt Ser	gaa Glu	gtg Val	atg Met	ctg Leu	tcc Ser	1056
			340						345						350	
act Thr	cgg Arg	att Ile	ggg Gly	tca Ser	ggc Gly	tct Ser	ttt Phe	gga Gly	act Thr	gtt Val	tat Tyr	aag Lys	ggg Gly	aaa Lys	tgg Trp	1104
			355						360						365	
cac His	gga Gly	gat Asp	gtt Val	gca Ala	gta Val	aag Lys	atc Ile	cta Leu	aag Lys	gtt Val	gtc Val	gac Asp	cca Pro	acc Thr	cca Pro	1152
370						375						380				
gag Glu	caa Gln	ttc Phe	cag Gln	gcc Ala	ttc Phe	agg Arg	aat Asn	gag Glu	gtg Val	gct Ala	gtt Val	ctg Leu	cgc Arg	aaa Lys	aca Thr	1200
385						390						395			400	

cgg	cat	gtg	aac	att	ctg	ctt	ttc	atg	ggg	tac	atg	aca	aag	gac	aac	1248
Arg	His	Val	Asn	Ile	Leu	Leu	Phe	Met	Gly	Tyr	Met	Thr	Lys	Asp	Asn	
				405					410					415		
ctg	gca	att	gtg	acc	cag	tgg	tgc	gag	ggc	agc	agc	ctc	tac	aaa	cac	1296
Leu	Ala	Ile	Val	Thr	Gln	Trp	Cys	Glu	Gly	Ser	Ser	Leu	Tyr	Lys	His	
				420					425					430		
ctg	cat	gtc	cag	gag	acc	aag	ttt	cag	atg	ttc	cag	cta	att	gac	att	1344
Leu	His	Val	Gln	Glu	Thr	Lys	Phe	Gln	Met	Phe	Gln	Leu	Ile	Asp	Ile	
				435					440					445		
gcc	cgg	cag	acg	gct	cag	gga	atg	gac	tat	ttg	cat	gca	aag	aac	atc	1392
Ala	Arg	Gln	Thr	Ala	Gln	Gly	Met	Asp	Tyr	Leu	His	Ala	Lys	Asn	Ile	
				450					455					460		
atc	cat	aga	gac	atg	aaa	tcc	aac	aat	ata	ttt	ctc	cat	gaa	ggc	tta	1440
Ile	His	Arg	Asp	Met	Lys	Ser	Asn	Asn	Ile	Phe	Leu	His	Glu	Gly	Leu	
				465					470					475		
aca	gtg	aaa	att	gga	gat	ttt	ggt	ttg	gca	aca	gta	aag	tca	cgc	tgg	1488
Thr	Val	Lys	Ile	Gly	Asp	Phe	Gly	Leu	Ala	Thr	Val	Lys	Ser	Arg	Trp	
				485					490					495		
agt	ggt	tct	cag	cag	gtt	gaa	caa	cct	act	ggc	tct	gtc	ctc	tgg	atg	1536
Ser	Gly	Ser	Gln	Gln	Val	Glu	Gln	Pro	Thr	Gly	Ser	Val	Leu	Trp	Met	
				500					505					510		
gcc	cca	gag	gtg	atc	cga	atg	cag	gat	aac	aac	cca	ttc	agt	ttc	cag	1584
Ala	Pro	Glu	Val	Ile	Arg	Met	Gln	Asp	Asn	Asn	Pro	Phe	Ser	Phe	Gln	
				515					520					525		
tcg	gat	gtc	tac	tcc	tat	ggc	atc	gta	ttg	tat	gaa	ctg	atg	acg	ggg	1632
Ser	Asp	Val	Tyr	Ser	Tyr	Gly	Ile	Val	Leu	Tyr	Glu	Leu	Met	Thr	Gly	
				530					535					540		
gag	ctt	cct	tat	tct	cac	atc	aac	aac	cga	gat	cag	atc	atc	ttc	atg	1680
Glu	Leu	Pro	Tyr	Ser	His	Ile	Asn	Asn	Arg	Asp	Gln	Ile	Ile	Phe	Met	
				545					550					555		
gtg	ggc	cga	gga	tat	gcc	tcc	cca	gat	ctt	agt	aag	cta	tat	aag	aac	1728
Val	Gly	Arg	Gly	Tyr	Ala	Ser	Pro	Asp	Leu	Ser	Lys	Leu	Tyr	Lys	Asn	
				565					570					575		
tgc	ccc	aaa	gca	atg	aag	agg	ctg	gta	gct	gac	tgt	gtg	aag	aaa	gta	1776
Cys	Pro	Lys	Ala	Met	Lys	Arg	Leu	Val	Ala	Asp	Cys	Val	Lys	Lys	Val	
				580					585					590		
aag	gaa	gag	agg	cct	ctt	ttt	ccc	cag	atc	ctg	tct	tcc	att	gag	ctg	1824
Lys	Glu	Glu	Arg	Pro	Leu	Phe	Pro	Gln	Ile	Leu	Ser	Ser	Ile	Glu	Leu	

```
<210> 7
<211> 668
<212> PRT
<213> Artificial Sequence
```

```

<400> 7
Met Glu His Ile Gln Gly Ala Trp Lys Thr Ile Ser Asn Gly Phe Gly
  1             5             10             15
Phe Lys Asp Ala Val Phe Asp Gly Ser Ser Cys Ile Ser Pro Thr Ile
      20             25             30
Val Gln Gln Phe Gly Tyr Gln Arg Arg Ala Ser Asp Asp Gly Lys Leu
      35             40             45
Thr Asp Pro Ser Lys Thr Ser Asn Thr Ile Arg Val Phe Leu Pro Asn
      50             55             60
Lys Gln Arg Thr Val Val Asn Val Arg Asn Gly Met Ser Leu His Asp
      65             70             75             80
Cys Leu Met Lys Ala Leu Lys Val Arg Gly Leu Gln Pro Glu Cys Cys
      85             90             95
Ala Val Phe Arg Leu Leu His Glu His Lys Gly Lys Lys Ala Arg Leu
      100            105            110
Asp Trp Asn Thr Asp Ala Ala Ser Leu Ile Gly Glu Glu Leu Gln Val
      115            120            125
Asp Phe Leu Asp His Val Pro Leu Thr Thr His Asn Phe Ala Arg Lys
      130            135            140

```


Arg His Val Asn Ile Leu Leu Phe Met Gly Tyr Met Thr Lys Asp Asn
405 410 415

Leu Ala Ile Val Thr Gln Trp Cys Glu Gly Ser Ser Leu Tyr Lys His
 420 425 430

Leu His Val Gln Glu Thr Lys Phe Gln Met Phe Gln Leu Ile Asp Ile
 435 440 445

Ala Arg Gln Thr Ala Gln Gly Met Asp Tyr Leu His Ala Lys Asn Ile
 450 455 460

Ile His Arg Asp Met Lys Ser Asn Asn Ile Phe Leu His Glu Gly Leu
 465 470 475 480

Thr Val Lys Ile Gly Asp Phe Gly Leu Ala Thr Val Lys Ser Arg Trp
 485 490 495

Ser Gly Ser Gln Gln Val Glu Gln Pro Thr Gly Ser Val Leu Trp Met
 500 505 510

Ala Pro Glu Val Ile Arg Met Gln Asp Asn Asn Pro Phe Ser Phe Gln
 515 520 525

Ser Asp Val Tyr Ser Tyr Gly Ile Val Leu Tyr Glu Leu Met Thr Gly
 530 535 540

Glu Leu Pro Tyr Ser His Ile Asn Asn Arg Asp Gln Ile Ile Phe Met
 545 550 555 560

Val Gly Arg Gly Tyr Ala Ser Pro Asp Leu Ser Lys Leu Tyr Lys Asn
 565 570 575

Cys Pro Lys Ala Met Lys Arg Leu Val Ala Asp Cys Val Lys Lys Val
 580 585 590

Lys Glu Glu Arg Pro Leu Phe Pro Gln Ile Leu Ser Ser Ile Glu Leu
 595 600 605

Leu Gln His Ser Leu Pro Lys Ile Asn Arg Ser Ala Ser Glu Pro Ser
 610 615 620

Leu His Arg Ala Ala His Thr Glu Asp Ile Asn Ala Cys Thr Leu Thr
 625 630 635 640

Thr Ser Pro Arg Leu Pro Val Phe Tyr Ser Phe Leu Pro Phe Phe Phe
 645 650 655

Phe Phe Phe Ser Phe Cys Phe Thr Pro Ser Thr Phe
 660 665

001120-204359

PATENT APPLICATION DECLARATION AND POWER OF ATTORNEY

My residence, post office address, and citizenship are as stated next to my name in PART A on page 2 hereof.

I believe I am the original, first, and sole inventor (if only one name is listed) or an original, first, and joint inventor (if plural names are listed) of the subject matter which is claimed and for which a patent is sought on the invention entitled **METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS USING TYROSINE KINASE RAF AND RAS**, the specification of which:

 X is attached hereto;

 was filed on as Application Serial No. and was amended
on (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to herein.

I acknowledge the duty to disclose information to the Patent and Trademark Office known to me to be material to the patentability of this application, as defined in Title 37, Code of Federal Regulations, Sec. 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Sec. 119 of any foreign application(s) for patent or inventor's certificate listed in PART B on page 2 hereof and have also identified in PART B on page 2 hereof any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

I hereby claim the benefit under Title 35, United States Code, Sec. 120 of any United States application(s) listed in PART C of page 2 hereof and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 25, United States Code, Sec. 112, I acknowledge the duty to disclose all information to the Patent and Trademark Office known to me to be material to patentability of this application, as defined in Title 37, Code of Federal Regulations, Sec. 1.56, which became available between the filing date of the prior application and the national or PCT international filing date of this application.

I hereby declare that all statements made herein of my knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

I hereby appoint the following as my attorneys or agents with full power of substitution to prosecute this application and transact all business in the United States Patent and Trademark Office connected therewith:

Michael A. Hierl	Reg. No. 29,807	Arne M. Olson	Reg. No. 30,203
Dolores T. Kenney	Reg. No. 31,269	Talivaldis Cepuritis	Reg. No. 20,818
Seymour Rothstein	Reg. No. 19,369	Daniel J. Deneufbourg	Reg. No. 33,675
Richard L. Robinson	Reg. No. 31,415	Joseph M. Kuo	Reg. No. 38,943
Mark Chao	Reg. No. 37,293	Martin J. Corn	Reg. No. 35,847

Whose mailing address for this application is:

OLSON & HIERL, LTD.
20 North Wacker Drive - 36th Floor
Chicago, Illinois 60606
Telephone: (312) 580-1180

PATENT APPLICATION DECLARATION AND POWER OF ATTORNEY

PART A: Inventor Information and Signature

Full name of SOLE or FIRST inventor John Hood
 Citizenship U.S. Residence 532.5J Via de la Valle
Solana Beach, California 92075
 Post Office Address (If different) _____

Inventor's signature: _____ Date: _____

Full name of SECOND Inventor Brian Eliceiri
 Citizenship U.S. Residence 3104 Hataca Road
Carlsbad, California 92009
 Post Office Address (If different) _____

Inventor's signature: _____ Date: _____

Full name of THIRD Inventor David A. Cheresch
 Citizenship U.S. Residence 3277 Lone Hill Lane
Encinitas, California 92024
 Post Office Address (If different) _____

Inventor's signature: _____ Date: _____


PART B: Prior Foreign Application(s)

<u>Serial No.</u>	<u>Country</u>	<u>Day/Month/Year Filed</u>	<u>Priority Claimed</u>
-------------------	----------------	-----------------------------	-------------------------

PART C: Claim for Benefit of Filing Date of Earlier U.S. Application(s)

<u>Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
60/148,924	August 13, 1999	
60/215,951	July 5, 2000	

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☒ Scanned copy is best available. *Drawings.*